

KK

SPIS RZECZY — CONTENTS

Nr 5

- Z. STEBNICKA. *Scarabaeoidea (Coleoptera)* of the Democratic People's Republic of Korea — *Scarabaeoidea (Coleoptera)* Koreańskiej Republiki Ludowo-Demokratycznej 191

Nr 6

- Z. SZYNDLAR. Herpetofauna Bieszczadów Zachodnich — The Herpetofauna of the Western Bieszczady Mts 299

Nr 7

- J. KOTEJA. Revision of the Genus *Exaeretopus* NEWSTEAD (*Homoptera, Coccidae*) — Rewizja rodzaju *Exaeretopus* NEWSTEAD (*Homoptera, Coccidae*). 337

POLSKA AKADEMIA NAUK
ZAKŁAD ZOOLOGII SYSTEMATYCZNEJ I DOŚWIADCZALNEJ

ACTA ZOOLOGICA CRACOVIENSIA

XXIV: 5—7

PAŃSTWOWE WYDAWNICTWO NAUKOWE
WARSZAWA—KRAKÓW 1980

RADA REDAKCYJNA — EDITORIAL BOARD

Przewodniczący — President: prof. dr R. J. Wojtusiak

Zast. przewodniczącego — Vice-President: doc. dr W. Micherdziński

Sekretarz — Secretary: dr hab. L. Sych

Członkowie — Members: doc. dr Z. Bocheński, prof. dr K. Kowalski, prof. dr M. S. Klimaszewski, prof. dr M. Młynarski, prof. dr J. Pawłowski, prof. dr J. Rafalski, prof. dr J. Razowski, prof. dr A. Riedel, prof. dr H. Szarski, prof. dr W. Szymczakowski

REDAKCJA — EDITORIAL STAFF

Redaktor naczelny — Editor-in-chief: prof. dr K. Kowalski

Zast. redaktora naczelnego — Subeditor: doc. dr Z. Bocheński,

Sekretarz — Secretary: dr hab. L. Sych

Adres redakcji: Zakład Zoologii Systematycznej i Doświadczalnej Polskiej Akademii Nauk,
ul. Sławkowska 17, 31-016 Kraków

Address of the Editor: Institute of Systematic and Experimental Zoology, Polish Academy
of Sciences, Sławkowska 17, 31-016 Kraków, Poland

Redaktor PWN
Maria Kaniowa

© Copyright by Państwowe Wydawnictwo Naukowe, Warszawa—Kraków 1980

ISBN 83-01-02098-9

ISSN 0065-1710

PAŃSTWOWE WYDAWNICTWO NAUKOWE — ODDZIAŁ W KRAKOWIE

Wydanie I. Nakład 810 + 90. Ark. wyd. 14,75. Ark. druk. 11²/₁₆ + 2 wklejki. Papier ilustr. kl. III 70×100 80 g.
Oddano do składania w grudniu 1979 r. Podpisano do druku w maju 1980 r. Druk ukończono w maju 1980 r.
Zam. 25/80 H-18 Cena zł 48,—

DRUKARNIA UNIwersytetu Jagiellońskiego w Krakowie

Zdzisława STEBNICKA

Scarabaeoidea (Coleoptera) of the Democratic People's Republic of Korea

[With 232 text-figs]

Scarabaeoidea (Coleoptera) Koreańskiej Republiki Ludowo-Demokratycznej

Abstract: The author presents a monographic elaboration of the *Scarabaeoidea* of the Democratic People's Republic of Korea. The systematic part of the paper contains a review of 138 species, including redescrptions of five species and description of one new to science. Extensive distributional data and locality records are presented as well as feeding habits and behaviour, where known. Results of a detailed zoogeographical analysis concerning the fauna of *Scarabaeoidea* of the whole Far East, complete the elaboration.

INTRODUCTION

This paper presents a first systematic and faunistic revision of *Scarabaeoidea* of the Democratic People's Republic of Korea (D.P.R.K.). It refers to the bibliographical data which concern the whole Korean Peninsula. The work has been accomplished basing on the materials collected by the participants of six zoological expeditions of Polish Academy of Sciences, who carried out investigations in the Democratic People's Republic of Korea in the years 1959—1974. A great number of specimens has been collected herself by the author of the present study during the 6th „Cracovian” expedition in 1974. Some specimens from Hungarian Museum of Natural History in Budapest and Zoological Institute of the Academy of Sciences of USSR in Leningrad, have also been examined. The systematic review of species with illustrations of their morphological features and redescrptions of the little known or rare species are contained in the systematic part of this study. The discussion of the particular species contains the references from systematic literature, significant titles from faunistic literature and bionomical and distributional data. Also author's own comments and the localities of investigated individuals have been given. Of the total number of 224 species known from the Korean Peninsula, 138 are discussed in the present study. This number contains 29 species which are new to the Korean fauna and one new to science. All the species are listed in Tab. II which also indicates

their general distribution. The localities in Korea mentioned in the text (including the localities taken from the cited literature) are numbered and marked on the schematic map of the Korean Peninsula (Fig. 3) according to index. The geographic names, which are given in the literature in the Japanese language, have been transcribed with the international transliteration applying latin alphabet (MROCZKOWSKI, 1972). Spelling of the majority of the names in the latin transliteration has been included in the index. The identification of North-Chinese and Manchurian localities that are indicated in the Japanese literature was not possible due to lack of the concerning cartographic sources. Following main works have been used for transliteration and identification of localities: The Columbia Lippincott Gazetteer of the World, Columbia, 1952; Bolšoj Korejsko-Russkij Slovar, Moscow, 1976; The World Atlas, Moscow, 1967; Atlas of the World, Warszawa, 1962.

Names of the institutions, which the examined specimens were from, are indicated in the text by the abbreviations listed below:

HMNH — Hungarian Museum of Natural History, Budapest

ISEZ — Institute of Systematic and Experimental Zoology of the Polish Academy of Sciences, Kraków

ZIL — Zoological Institute of Academy of Sciences of USSR, Leningrad

ZIW — Zoological Institute of the Polish Academy of Sciences, Warszawa.

The author wishes to express her cordial thanks to the following persons for their valuable help and cooperation: To Dr. hab. M. MROCZKOWSKI from the Zoological Institute of the Polish Academy of Sciences in Warszawa for lending use of the materials collected during previous expeditions and for introduction to his own observations; to Dr. S. ENDRÖDI from Budapest and Dr. O. L. KRYZHANOVSKIJ from Leningrad for their kind making available the comparative materials; to Mr. ČHOI-DE-VŎN and Mr. KIM-HJÖNG-SÖK, the functionaries of the Korean Academy of Sciences, and also to Mr. DŽU-DONG-JUL, Director of the Zoological Institute for their help in the organization of the 6th expedition and contribution in the course of it; to Prof. J. PAWŁOWSKI and Dr. hab. A. SZEPTYCKI, the participants of expedition for their interest and involvement in collecting *Scarabaeoidea*.

HISTORICAL REMARKS ON THE INVESTIGATIONS

Examination of *Scarabaeoidea* of the Korean Peninsula was undertaken already in the 19th century, however, in the European and Japanese systematic-faunistic literature only short notes concerning single species have appeared. The ditributional data were usually limited to a single word „Korea”; in some cases totally disfigured names of provinces or larger towns were exclusively available. Only a few materials were collected coincidentally (mainly by the investigators who were interested in other animal groups) and therefore they are not accompanied by informations concerning the environment of the col-

lected specimens. Up to now, a Japanese coleopterologist Jozo MURAYAMA was the only author who had published in a separate work the results of his many years' investigations on *Scarabaeidae pleurosticti* of Manchuria and Korea. He spent some time at the state agencies in these countries, which were Japanese provinces then. For 24 years he has collected the material and subsequently has published small reports (MURAYAMA, 1931; 1934; 1937; 1938; 1941). Simultaneously he was preparing the edition of a monography. The story of this book is very dramatic: in the war the manuscript and the notes have been buried. The reconstruction of them took some years and finally the book was published in Japan at „Aid of the Department of Education” in 1954. Unfortunately, before his death J. MURAYAMA managed to publish only the first volume of the monography, which covers only part of the collected materials, namely 74 species of *Melolonthinae*. The rest of the MURAYAMA's collection is probably in China now. His book has been published in Japanese language; it contains descriptions of new species, redescriptions of the known ones, drawings of the male copulatory apparatuses and color pictures of beetles on which the main morphologic features are emphasized. A list of references to the literature and detailed localities of the species in Manchuria and Korea, also in the southern part of the Korean Peninsula are given. This work is of an extraordinary relevance; it helps to solve many taxonomic problems, since it deals with many local species, which in the previous papers were discussed very superficially.

In the nineteen fifties the Polish Academy of Sciences has undertaken very intensive investigations on the fauna of the northern part of the Peninsula. The arrangement concerning scientific collaboration has been established between us and the Academy of Sciences of the Democratic People's Republic of Korea. Also the Hungarian Museum of Natural History joined the investigations and organized two zoological expeditions in the years 1970—1971. The participants of six Polish zoological expeditions in the years 1959, 1965, 1966, 1970, 1971, 1974 gathered a huge collection of invertebrates, which resulted in numerous papers by Polish and European specialists of various systematic groups. As it has been mentioned at the beginning, the present paper, which is a partial review of the investigations done up to the present, was accomplished thanks to a very abundant collection gathered by the Polish expeditions.

ZOOGEOGRAPHY

1. Methodical part

The analysis of the Korean fauna of *Scarabaeoidea* and its connections with the faunas of the contiguous territories is the subject of the present discuss. It has been done based on the present knowledge of these faunas. The tables




Fig. 1. Diagram of affinities between the faunas of compared territories, A — Taiwan; B — Ryu-kyu, Izu Archipelagos; C — Honshu, Shikoku, Kyushu; D — Hokkaido, Sachalin, Kurile Islands; E — Korean Peninsula; F — Manchuria, Priamurie; G — Mongolia; H — North China; I — Central China; K — South China; L — Indochina. Numbers mark the percentile affinity factors

given include 224 species of the Korean Peninsula recorded in the previous literature as well as new species, which are discussed in the systematic part of the present paper. The tables also contain the data concerning general distribution of particular species; some geographic territories bordered on the schematic map (Fig. 2) and marked with letters were considered. The distribution of the species which occur in the distinguished and compared regions of the Eastern Asia, was obtained from the data contained in the catalogues, monographies and numerous faunistic papers. The affinity criterium has been applied to acquire an objective idea of the similarities among the faunas in the distinguished regions. The percentile version of the SØRENSEN'S formula (1948)

was applied to the statistical analysis: $S = 100 \times \frac{2c}{a+b}$, in which S = affinity factor, a, b = number of species in the compared geographic territories, c = number of common species. The faunistic affinity factors obtained are illustrated in the diagram (Fig. 1), which has been plotted according to CZEKANOWSKI (1930). This diagram confirms in general that the classification of the compared regions has been done properly. However, it is not ideal, since as a principle the greatest similarities should be located as close to the diagonal line of the diagram as possible. Nevertheless, fairly objective picture of the differentiation of Scarabaeoidean fauna of this part of Far East was acquired (Fig. 2). The obtained results indicate the importance of the ecological, environmental and historical factors, which have an effect on the systematic group under discussion; they are considered further on.

2. Zoogeographic analysis

The superfamily *Scarabaeoidea* is not uniform as to the role which its elements play in the environment. What is more, both its systematics and chorology are not totally understood. This especially concerns the group of subfamilies *Scarabaeidae laparosticti* which, in contradistinction to the group of *S. pleurosticti* contains species that are not linked to the host plants, have different ecologic preferences and indicate adaptative mechanisms much more difficult to detect. For the above reasons common discussion of both groups could affect the value of the obtained results, which determine not only the affinity of faunas among particular territories, but also indirectly shows the similarity of the landscapes and environments which the discussed faunas are linked to.

In order to illustrate the comparisons of Korean fauna with the faunas of the distinguished territories, the affinity factors for each group separately as well as for both groups together have been calculated and given in the table below:

Table 1

E. Korea Adjacent territories	Affinity factors (%)		
	<i>S.</i> <i>laparost.</i>	<i>S.</i> <i>pleurost.</i>	<i>Scarabaeoi-</i> <i>dea</i>
A. Taiwan	17	17	17
B. Ryu-kyu, Izu Isl.	31	23	26
C. Honshu, Shikoku, Kyushu	46	34	39
D. Hokkaido, Sachalin, Kurile Isl.	44	29	34
F. Manchuria, Priamurie	46	58	53
G. Mongolia	27	19	22
H. North China	42	38	41
I. Central China	19	28	24
K. South China	11	10	10
L. Indochina	7	2	4

The above specification indicates that the affinity factor in the majority of regions compared to Korea is higher in the group of *S. laparosticti* than in the group of *S. pleurosticti*; these differences due to the following reasons. The group of *S. pleurosticti* is in general better known faunistically. Therefore the number of species given in this group is relatively high. The majority of them are, however, local species whose area of distribution is sometimes very narrow. Consequently, the composition of species is different in the compared regions and the number of common species is strikingly low in relation to the total fauna. There is an opposite situation in *S. laparosticti*; the number of endemic species is considerably lower, while numerous given species have very wide distribution. Therefore the number of common elements is higher and the percentage of the faunistic affinity is consequently increased. To some extent, the indicated differences also determine the environmental factors which affect each of the groups, however, detailed discussion of this problem seems untimely. Further faunistic investigations, in particular those carried out in the little known regions as, e. g. China, shall probably change the numerical relationships between both groups. It seems, however, that the general idea of the affinities that join all the *Scarabaeoidea* in the compared regions shall remain unchanged. The numerical data concerning the whole superfamily are given in the diagram (Fig. 1). They illustrate some regularities in the geographic distribution of *Scarabaeoidea* in quite a convincing way. The distribution of this superfamily is fairly similar to the distribution of other groups of entomofauna in this part of Palaearctic and Oriental Realm.

The analysis of the diagram (Fig. 1) leads to the following conclusions for the particular regions:

A. Fauna of Taiwan, which is considered by some authors (KOSTROWICKI, 1965) as a separate region of the Oriental Realm, is characterized by relatively low affinity factors (maximum 23%). This fact indicates a great independence of this fauna, which contains numerous endemic species. Its closest

connections are with the fauna of Central China (23%), South China (20%), the fauna of contiguous islands of the archipelago Ryu-kyu (18%) and Korean Peninsula (17%).

B. Relatively poor fauna of the archipelagos Ryu-kyu and Izu contains few endemic species, some species of wider insular and continental distribution and the subspecies, that were distinguished by Japanese authors based on the variability of populations inhabiting particular islands. It corresponds to the fauna of the islands of Central Japan (35%), Korean Peninsula (26%), northern insular region that contains Hokkaido (23%) and to the fauna of Taiwan (18%). Taking into account the distribution of *Scarabaeoidea* one may consider the archipelago Ryu-kyu as a Palaearctic-Oriental transitional zone of the insular sector. This zone is located more or less between the islands Yakushima and Amami-Ōshima („WATASE-line”, Fig. 2) and the islands Miyako and Ishigaki in the southern part of the archipelago („HACHISUKI-line”, Fig. 2).

C. The islands of the Central Japan include Honshu, Shikoku, Kyushu and Tsushima (the latter located very close to Korean Peninsula) as well as small neighbouring islands, among others Sado and Oki. Fauna of these islands is most closely related to the faunas of Hokkaido — Sachalin — Kurile Islands (50%), Korean Peninsula (39%), Ryu-kyu Archipelago (23%) and Manchurian-Priamurian fauna (22%). Less close similarities are those to the fauna of North and Central China (18% each). Scarabaeoidean fauna is fairly uniform there. Only in the northern part of Honshu, which extends beyond the subtropical zone, does a slightly different fauna occur; it contains northern elements. According to phytogeographers (GOOD, 1974) and zoogeographers (KURENCOV, 1961), the islands of the Central Japan and Korean Peninsula constitute Korean-Japanese phyto- and zoogeographic region; both flora and fauna of this region is genetically and historically closest to forest formations and to faunas of Manchurian-Priamurian and North-Chinese territories.

D. Fauna of the region that includes Hokkaido, Kurile Islands and Sachalin is closest to the faunas of Central Japan (50%), Korean Peninsula (34%), Manchuria-Priamurie (27%), and Ryu-kyu Islands (23%). It is also closely related to the fauna of Northern China (20%). The island of Hokkaido, which is separated from the northern promontory of Honshu by the narrow, geologically young straits of Tsugaru („BLAKISTON-line”, Fig. 2), contains high percent of Central-Japanese species. It also contains boreal elements and its faunistic composition corresponds to those of Manchurian-Priamurian and Korean territories. There is a very poor fauna in Sachalin and Kurile Islands. It is mainly composed of boreal-northern and Manchurian-Priamurian elements.

The numeral data discussed above indicate that the fauna of *Scarabaeoidea* in the regions of the insular sector is characterized by the closest reciprocal relations. Taiwan, the fauna of which is more closely linked to the continental one of Central and South China, is an exception.

E. Korean Peninsula is centrally located in the discussed regions. Thus, their faunas largely contribute to the composition of the Korean fauna; this

is reflected in fairly high values of the indicators. The ordered affinities are as follows: Manchuria-Priamurie (53%), North China (41%), Central Japan (39%), Hokkaido-Sachalin-Kurile Islands (34%), Ryu-kyu, Izu (26%), Central China (24%), Mongolia (22%). These are followed by the faunas of Taiwan (17%), South China (10%) and Indochina (4%).

There are five main types of fauna found in Korean Peninsula:

- a) Eurosiberian and Palaearctic species, which are nearly uninterruptedly distributed on the whole territory of Siberia and Europe;
- b) less and less common, vanishing in the direction of West, species that inhabit the forests and open areas in Manchuria and South-East Siberia;
- c) East-Asiatic species widely distributed in Manchuria and South-East Siberia, Japan and China; they form the most numerous group;
- d) species that constitute the local fauna of Korean Peninsula;
- e) Oriental species, which are widely distributed in South and Central China, Taiwan and southern Japanese Islands, as well as in Indochinese Peninsula and Philippines.

F. Manchuria, South-Eastern Siberia (Amur-Ussuri Lands and Primorskiy Kraj = „Priamurie”). Most complex is the fauna of mixed and leafy forests, described by zoogeographers as „Manchurian-type”. It inhabits a separate zoogeographic subregion, whose borders determined by WALLACE (1876) are perpetually changed. Botanists (KOMAROV, 1949; ALECHIN, 1951) have largely contributed to the knowledge of the term „Manchurian elements”. They determined the area of Manchurian flora beyond Manchuria down to Amur—Ussuri Lands and North Korea. In the recent years due to the investigation of the distributional areas of various systematic groups of East-Asiatic entomofauna it was discovered, that its main part was concentrated in this very region. It is closely linked to the floristic landscape and is characterized by the occurrence of both endemic species and endemic genera. The fauna of *Scarabaeoidea*, in particular of the group of *S. pleurosticti*, is extremely rich in this region. In the north it borders upon a very poor tundra-taiga (Ochotsk—Kamchatka) fauna, in the west upon Mongolian-Daurian steppe fauna, in the south and south-east upon the North-Chinese and Korean-Japanese faunas connected to plant formations in which the evergreen plants are contained. The highest affinity factors indicate the connections of this fauna with those of Korea (53%), North China (45%), Mongolia (28%), Hokkaido-Sachalin-Kurile Islands (27%) and Central Japan (22%). Its connections with the fauna of Central China are insignificant (13%).

G. Mongolia is one of the regions that are better known faunistically. By zoogeographers (KOSTROWICKI, 1965) it is included into the Central-Asiatic Province, which contains the faunas of the forests and wooded-steppes of the Central Asiatic mountains and also the fauna of dry steppes, half-deserts and deserts of the central part of Palaearctic. Relatively low values of the factors indicate that the Mongolian fauna is separated zoogeographically from the

other discussed regions. It is closest to the faunas of Manchuria-Priamurie (28%), North China (25%) and Korea (22%).

H. North-Chinese territory reaches Huanghe in the south and more or less Sinkiang-region in the west. This region is very little known; based on the present knowledge of the fauna of *Scarabaeoidea*, the highest affinity factors indicate its connections with those of Manchuria-Priamurie (45%), Korean Peninsula (41%), Mongolia (25%) and Central China (24%). Less close similarities are those to the fauna of Hokkaido-Sachalin-Kurile Islands (20%) and Central Japan (18%).

I. Central China extends between Huanghe and Yangzi (Changjiang), in the west it reaches Sichuan excluding Tibet. Kansu, Sichuan Valley and the eastern part which borders upon the Yellow Sea are partially better investigated. Considerable separateness is indicated by the fauna of Sichuan Valley and surrounding mountains, however, precise comparison are not able since the composition of species in Central China is poorly known and exact indicators are lacking. The closest faunas are those of South China (33%), North China and Korean Peninsula (24% each) and of Taiwan (23%). Connections with the fauna of Central Japan are also marked (18%), while the affinity to the Palaearctic-Oriental fauna of Ryu-kyu is very weak (10%). Based on the present knowledge of distribution, the eastern part of the discussed region, which is located between the rivers Huanghe, Yangzi and Weihe, should be recognized as Palaearctic-Oriental transitional zone.

K. The South-Chinese territory, which belongs to Oriental Realm, is situated between the river-basin of Yangzi and the borders of Indochinese Peninsula (Birma, Vietnam); it also includes the island Hainan. This area is little known faunistically. The majority of reports concern the provinces of Yunnan and Fukien as well as Hainan Island, therefore the indicators are not fully reliable. The sequence of affinities with the neighbouring faunas is as follows: Central China (33%), Indochina (24%), Taiwan (20%) and Korean Peninsula (10%). The rest of the factors do not exceed 8% (Ryu-kyu).

L. The Indochinese territory encircled on the map (Fig. 2) has been treated as the example of the faunistic separateness of the typically Oriental zone. It indicates very weak connections with the palaearctic part of Eastern Asia (the indicators do not exceed 4%). Unlike China, this area is better known. Thus it seems that the affinity factor of the closest fauna of South China has a lowered value (24%). More loose connections of this fauna have been found with the faunas of Taiwan (14%) and Central China (11%).

The results of the investigations done up to the present indicate that the fauna of *Scarabaeoidea* in Palaearctic and Palaearctic-Oriental part of Eastern Asia is very complex. It consist of groups that are historically connected to the evolution of the landscape. Depending on the geological basis and on the isolation from the neighbouring forms they indicate autochthony and endemicity of various degrees. The fauna of Korea is one of such complexes; it

has been formed for a very long time. It is particularly interesting because of the geomorphologic features of the territory which is linked to. Unsatisfactory knowledge of the fauna in the southern part of Korean Peninsula, mainly of




Fig. 2. Division of the studied area into working territories with diagram of affinities of the Korean fauna

its most southern coastal regions, makes the formulation of conclusions extremely difficult. A detailed examination of the whole Korea with contiguous islands would certainly result in many new zoogeographic observations. This is particularly important because the Korean fauna undergoes further transformation, connected among others to human activity.

The reconstruction of the history of the present fauna of the Far East is based on the palaeogeography and history of flora, which were the subject of several studies (ALECHIN, 1951; KOSTROWICKI, 1969; KRIŠTOFOVIČ, 1932, 1958; KURENCOV, 1952, 1960, 1961, 1974; SOLOVEV, 1961). The performance of the history of this fauna is very difficult, since various faunistic groups as, e. g., faunas of leafy forests of the European type, of the coniferous forests, faunas of the open areas and evergreen forests, were formed as soon as in the Tertiary and shifted multidirectionally due to succeeding changes of the landscape and climate.

The marine regression caused by the intense tectonic movements, lasted during almost the entire Palaeogene. The Japanese Sea, which in the Palaeocene and Eocene was only a shallow gulf, completely disappeared in Oligocene, thus nearly whole Japan became connected with the continent. This region was under influence of temperate-warm and wet climate. Hence it was overgrown with the forest of Greenland type (domination of deciduous leafy trees and numerous conifers), only in the south were there mixed forests of the Greenland-subtropical type (with participation of evergreen trees and bushes). The orographic situation at that time was similar to this at present: three landscape types dominated — mountainous (Sichote-Alin, North Korea and Japan), uplands (West China) and lowlands (East China, Amur, Ussuri Land). The faunas of the lowland plains and borders of the fresh-water lakes that expanded out of drying Mongolia were developing parallelly to the development of the forest fauna. Higher in the mountains, formation of an ancient nucleus of Alpine fauna probably took place. Subsequently, strong tectonic movements and volcanic activity considerably affected the configuration of the ground: the massives like Chingan in Manchuria and numerous mountain chains in Korea were formed. The denudated sediments from these mountains covered the whole Yellow Sea, which was land at that time. Due to the processes of Alpine orogenesis and formation of steppes and deserts of Central Asia, on the turn of Palaeogene and Neogene the border of the subtropical flora and fauna shifted to the north. The vegetation of continental China and Mongolia underwent gradual xerophilisation.

Xerophilous brushwoods appeared in Manchuria and Korea as well; they bore down the plants demanding more humidity. Mongolia became covered with xerophilous vegetation and its humidity-demanding fauna mortified with the recession of lakes and lowland plains, which were similar to the former ones near Lower Amur. By drawing off into east this fauna probably sheltered partially in slightly similar environments on the plains of lake Chanka and Amur.

In a short period in Miocene, Sachalin and Japan formed an archipelago of small islands. They were surrounded by a shallow sea, which at the turn of Miocene and Pliocene withdrew behind the line Japan-Kurile Islands-Kamchatka and consequently the islands totally joined the continent again. Probably due to this reason the climate of the islands, as well as of the continental coastal zones that were affected periodically by the pressure of the ocean, was rather cool and wet, while on the rest of the region, temperate-subtropical climate was dominated. The xerophilisation of flora attained the islands only partially. It did not result in impetuous and acute ecological differentiations; in the most favorable refuges the fauna indicated a tendency to conserve the ancient forms. On the contrary, the fauna of the western part of the region remained in less favorable conditions: steppe-desert fauna that originated from Central Asia, assimilated the Tertiary forest fauna of North China. This strongly affected the ecological differentiation of biocenoses in Manchuria and Korea.

The period of older glaciation (Riss) was characterized by cooling of the climate. This was connected to the development of the mountain glaciers, which were strongest in the North-East Siberia and occurred locally in the mountains of Hokkaido and Honshu, Sichote-Alin, Chingan and North Korea. In this period, the wooded-steppe type vegetation, including tundra-growing species, dominated in the north and steppe flora in the south and the west. Forest formations were only locally preserved in the south and in the mountains, among others in Korea. The early glaciation only caused an indirect change in natural conditions of the Far East, through general cooling of the climate and shifting of some groups of thermophilous species to the south. Through dislocation along the present Korean Peninsula these faunas could contact those of Japan and China again.

In the interglacial warm period the seas of Far East gradually appeared; this led to a renewed humidification of the continental climate. Forests dominated in the east and south again; they were slightly less abundant than in the pre-glacial period. On the other hand in the western part of region, large areas were covered with wooded-steppe, characterized by more abundant vegetation. Manchurian, Priamurian and Korean faunas maintained the connection with the Japanese fauna through a narrow stripe of land, which still occurred instead of the present Korean Straits. This concerned Sachalin as well, which was also connected with the continent then. Thus, the close faunal connections established a long time ago were still sustained.

The last glaciation (Würm) affected more distinctly the natural conditions of Far East and therefore caused considerable disturbances in the biocenoses. Forest-tundra dominated in the north, while in the south — open steppes of the Mongolian type. Forest vegetation, mostly coniferous, occurred locally in the mountains of Sichote-Alin, in South Chingan and in Korea. Some ecologic complexes of the contemporary fauna sheltered themselves in the southern refuges, among others in the south of Primorskij Kraj, in Korea and in the

south of Manchuria. There they formed sort of reservation; it was surrounded in the periphery by a zone of relicts that survived up to the present in the form of extremely impoverished complexes or single species adapted to the ecologic requirements of other faunas. Some elements of the thermophilous, subtropical fauna did not stand the cooling of the climate and perished. Therefore the northern border of the area of this fauna became shifted towards the south, into Central China. This was followed by the parallel succession of the fauna of tundra in the direction of the south. This was accomplished among others along the comb of Kurile Islands, which then served as a bridge to the large Japanese Islands.

Towards the close of Pleistocene, due to tectonic activity of the bottom of the sea, the Japanese Islands and Sachalin became finally separated from the continent. In the same period numerous straits and Kurile Islands were created in their present form; in the whole the Quaternary period the islands were affected by the tectonic movements and consequently their shape and surface configuration were not stable. The vegetation of the forest formations became dominant again, however, these forests were not nearly so rich as those of the early Pleistocene.

During the post-glacial warming up the faunas came back from the south accompanied by the plant formations. They established a biocenotic contact with the faunas of the northern, north-western and western refugia. The process of settlement of islands by the continental species was stopped (in the case of invertebrates was rather strongly limited) at the decline of Pleistocene. The migration pathways leading from the North-Eastern Asia through Sachalin — Hokkaido to Honshu and through Korea to southern part of Honshu were cut by the sea. Although there is no proof that the land bridge in the part of Honshu—Ryu-kyu—Taiwan did exist, the faunistic affinities of these islands support the assumption that the ways of expansion of species as, e. g., mechanical introduction due to typhoons or means of marine transport do occur.

The present period is marked by the limitation of forest grown area by man. In China forests have been almost completely felled over large areas and their habitat turned into farm land. On the huge territories contiguous to Amur there are vast meadows. In the Korean Peninsula large areas of the old forests were burned away with napalm in the war. Some enclaves of the old forests survived still locally in the Chingan mountains, Sichote-Alin and Korea. Most numerous Tertiary plant formations are maintained in Japan and Korea; they represent the refuges of the relicts of numerous faunistic groups.

The determination of which species or groups of species of *Scarabaeoidea* followed particular ways to the Korean Peninsula is very difficult because in the whole period of Tertiary and in main part of Quaternary, the discussed region was the scene of migrations both ways: from the depths of the continent to the present islands and backwards. In the period of Quaternary fluctuation of climate, the Korean pathway was possibly followed by the southern species, which in the warmer periods migrated into north, as well by the north-

No	Species	A	B	C	D	E	F	G	H	I	K	L	Eurasia
* 25	<i>Onthophagus (Onthophagus) rugulosus</i> HAR.	•				•				•	•	•	
* 26	<i>Onthophagus (Onthophagus) olsoufieffi</i> BOUC.					•	•		•				
* 27	<i>Onthophagus (Onthophagus) punctator</i> REITT.					•	•						
28	<i>Onthophagus (Onthophagus) necessarius</i> REITT.					•							
29	<i>Onthophagus (Onthophagus) pupillatus</i> KOLBE					•							
30	<i>Onthophagus (Onthophagus) trituber</i> (WIED.)	•				•				•	•	•	
31	<i>Onthophagus (Onthophagus) koma</i> MATS.					•							
* 32	<i>Aphodius (Colobopterius) apicalis</i> HAR.			•		•			•				
* 33	<i>Aphodius (Colobopterius) indagator</i> MANNRH.					•							
* 34	<i>Aphodius (Colobopterius) subterraneus</i> (L.)					•	•						•
* 35	<i>Aphodius (Teuchestes) brachysomus</i> SOLS.					•							
* 36	<i>Aphodius (Teuchestes) haemorrhoidalis</i> (L.)				•	•			•				•
* 37	<i>Aphodius (Teuchestes) donghariensis</i> STEB.					•							
* 38	<i>Aphodius (Pharaphodius) rugosostriatus</i> WATERH.					•			•				
* 39	<i>Aphodius (Coptochiroides) subcostatus</i> KOLBE					•			•				
* 40	<i>Aphodius (Acrossus) rufipes</i> (L.)	•	•		•	•	•						•
* 41	<i>Aphodius (Acrossus) binaevulus</i> HEYD.					•							
* 42	<i>Aphodius (Acrossus) superatratus</i> NOM. et NAK.					•							
* 43	<i>Aphodius (Paulianellus) maderi</i> BALTH.					•							
* 44	<i>Aphodius (Pleuraphodius) lewisi</i> WATERH.	•				•					•		
* 45	<i>Aphodius (Trichaphodius) proclivis</i> BALTH.		•			•					•		
* 46	<i>Aphodius (Trichaphodius) comatus</i> A. SCHM.					•			•				
* 47	<i>Aphodius (Aganacrossus) urostigma</i> HAR.	•	•			•					•		
* 48	<i>Aphodius (Pseudacrossus) justus</i> PETR.					•							
* 49	<i>Aphodius (Trichonotulus) dzamosanicus</i> STEB.					•							
* 50	<i>Aphodius (Trichonotulus) mongolicus</i> MANNRH.					•							
* 51	<i>Aphodius (Aphodius) elegans</i> ALLIB.	•	•			•					•		•
* 52	<i>Aphodius (Orodatus) pusillus</i> (HBST)					•							
* 53	<i>Aphodius (Orodatus) naraensis</i> NAK.					•							
54	<i>Aphodius (Aphodaulacus) nigrotessellatus</i> MOTSCH.					•							
55	<i>Aphodius (Volinus) obsoleteguttatus</i> WATERH.					•							
* 56	<i>Aphodius (Phaeaphodius) rectus</i> MOTSCH.	•	•			•							
* 57	<i>Aphodius (Agrilinus) uniformis</i> WATERH.					•							
* 58	<i>Aphodius (Agritinus) putridus</i> (HBST)					•							
* 59	<i>Aphodius (Agrilinus) brevisculus</i> (MOTSCH.)	•				•							•

No	Species	A	B	C	D	E	F	G	H	I	K	L	Eurasia
* 60	<i>Aphodius (Agrilinus) inexpectatus</i> BALTH.					•	•						
* 61	<i>Aphodius (Bodilus) sordidus</i> (FABR.)		•	•	•	•	•		•	•	•		•
* 62	<i>Aphodius (Bodilus) languidulus</i> A. SCHM.					•	•		•	•			
* 63	<i>Aphodius (Niatius) sturmi</i> HAR.			•		•		•					•
* 64	<i>Aphodius (Calamosternus) uniplagiatus</i> WATERH.		•	•	•	•	•						
* 65	<i>Aphodius (Calamosternus) sublimbatus</i> MOTSCH.		•	•	•	•	•		•	•	•		
66	<i>Aphodius (Calamosternus) prusai</i> TES.					•	•						
67	<i>Aphodius (Pharaphodius) orientalis</i> HAR.		•	•		•	•			•	•		
* 68	<i>Aegialia (Aegialia) comis</i> (LEWIS)			•	•	•	•					•	
* 69	<i>Aegialia (Psammoporus) kamtschatica</i> MOTSCH.			•	•	•	•						
* 70	<i>Rhyssenus koreanus</i> sp. n.					•	•						
71	<i>Saprosites japonicus</i> WATERH.		•	•	•	•	•						
* 72	<i>Ochodaeus ferrugineus</i> (ESCH.)					•	•					•	
73	<i>Kolbeus koreanus</i> KOLBE		•	•		•	•			•			•
* 74	<i>Geotrupes (Phelotrupes) auratus</i> MOTSCH.					•	•						
* 75	<i>Geotrupes (Phelotrupes) laevistriatus</i> MOTSCH.		•	•	•	•	•		•	•			
76	<i>Bolbocerodema zonatum</i> NIKOL.					•	•		•				
77	<i>Bolbocerosoma nigroplagiatum</i> (WATERH.)		•	•	•	•	•						
* 78	<i>Gastroserica herzi</i> (HEYD.)	•	•			•	•			•			
* 79	<i>Nipponoserica similis</i> (LEWIS)			•		•	•						
* 80	<i>Trichoserica polita</i> (GEBL.)					•	•						
* 81	<i>Sericania fuscilineata</i> MOTSCH.		•	•		•	•						
82	<i>Sericania latiusculata</i> MURAY.					•	•						
83	<i>Sericania koryoensis</i> MURAY.					•	•						
84	<i>Sericania hasegawai</i> MURAY.					•	•						
85	<i>Paraserica grisea</i> (MOTSCH.)			•	•	•	•						
* 86	<i>Serica boops</i> WATERH.		•	•	•	•	•						
87	<i>Serica brunnea</i> (L.)			•	•	•	•						•
88	<i>Serica septentrionalis</i> MURAY.					•	•						
89	<i>Serica elliptica</i> MURAY.					•	•						
* 90	<i>Maladera (Maladera) castanea</i> (ARR.)	•	•	•	•	•	•		•	•			
* 91	<i>Maladera (Maladera) gibbiventris</i> (BRSKE)	•	•	•	•	•	•						•
* 92	<i>Maladera (Maladera) formosae</i> (BRSKE)	•	•	•	•	•	•						•
* 93	<i>Maladera (Maladera) renardi</i> BALL.	•	•	•	•	•	•		•				•

No	Species	A	B	C	D	E	F	G	H	I	K	L	Eurasia
* 94	<i>Maladera (Maladera) holosericea</i> (SCOP.)					•	•						•
* 95	<i>Maladera (Maladera) cariniceps</i> (MOS.)	•		•		•	•			•			
* 96	<i>Maladera (Maladera) laboriosa</i> (BRSKE)	•				•							
* 97	<i>Maladera (Maladera) fusaria</i> MURAY.	•				•							
* 98	<i>Maladera (Maladera) orientalis</i> (MOTSCH.)	•	•	•	•	•	•	•					
* 99	<i>Maladera (Maladera) schonfeldti</i> (MURAY.)					•							
100	<i>Maladera (Maladera) opaciventris</i> (MOS.)					•							
101	<i>Maladera (Maladera) thibetana</i> (BRSKE)					•							
* 102	<i>Maladera (Maladera) okamotoi</i> (MURAY.)					•							
103	<i>Maladera (Maladera) infuscata</i> (MOS.)	•				•				•			
104	<i>Maladera (Maladera) stridula</i> (BRSKE.)	•				•				•			
105	<i>Maladera (Maladera) japonica</i> (MOTSCH.)	•	•	•	•	•							
106	<i>Maladera (Maladera) verticalis</i> (FAIRM.)	•				•	•	•	•				
107	<i>Maladera (Maladera) ovatula</i> (FAIRM.)					•	•	•	•				
108	<i>Maladera (Maladera) aureola</i> (MURAY.)					•							
109	<i>Maladera (Maladera) koreana</i> (MURAY.)					•							
* 110	<i>Maladera (Eumaladera) nitidiceps</i> NOM.		•			•							
* 111	<i>Hoplia (Euchromoplia) aureola</i> (PALL.)					•		•	•				
* 112	<i>Hoplia (Decamera) djukini</i> JACOB.					•							
* 113	<i>Ectinohoplia rufipes</i> (MOTSCH.)			•	•	•			•	•			
114	<i>Polyphylla laticollis</i> LEWIS			•		•							
115	<i>Polyphylla chinensis manchurica</i> SEM.					•							
* 116	<i>Hoplosternus incanus</i> MOTSCH.			•		•			•	•			
117	<i>Hoplosternus japonicus</i> HAR.			•		•			•				
* 118	<i>Hylotrogus bicoloratus</i> (HEYD.)					•							
119	<i>Lachnosterna nipponensis</i> MURAY.			•	•	•							
120	<i>Heptophylla picea</i> MOTSCH.			•		•							
* 121	<i>Apogonia cupreoviridis</i> KOLBE			•		•							
* 122	<i>Apogonia cribricollis</i> BURM.			•		•							
123	<i>Cyphochilus farinosus</i> WATERH.			•		•			•				
* 124	<i>Brahmina rubetra</i> (FALD.)			•		•			•				
125	<i>Brahmina intermedia</i> (MANNRH.)					•			•				
126	<i>Brahmina darcisi</i> REITT.					•							
127	<i>Brahmina crenicollis</i> MOTSCH.					•							

No	Species	A	B	C	D	E	F	G	H	I	K	L	Eurasia
128	<i>Brahmina excisiceps</i> MURAY.					•				•			
* 129	<i>Lasiopsis manchuricus</i> MURAY.					•	•						
130	<i>Lasiopsis sahbergi</i> (MANNRH.)					•	•						
131	<i>Metabolus impressifrons</i> FAIRM.					•			•				
* 132	<i>Sophrops heydeni</i> (BRsKE)					•							
* 133	<i>Sophrops striata</i> (BRsKE)					•							
* 134	<i>Miridiba koreana</i> NIJ. et KIN.					•			•				
135	<i>Miridiba castanea</i> (WATERH.)			•		•							
* 136	<i>Holotrichia diomphalia</i> (BAT.)			•		•			•				
* 137	<i>Holotrichia ernesti</i> REITT.				•	•			•				
* 138	<i>Holotrichia paralella</i> (MOTSCH.)			•		•			•				
* 139	<i>Holotrichia picea</i> WATERH.				•	•			•				
* 140	<i>Holotrichia oblita</i> (FALD.)			•		•							
* 141	<i>Holotrichia inelegans</i> (LEWIS)			•		•							
142	<i>Holotrichia castanea</i> (WATERH.)			•		•							
143	<i>Holotrichia sichotana</i> (BRsKE)					•			•				
144	<i>Holotrichia kiotoensis</i> (BRsKE)			•		•			•				
145	<i>Holotrichia reticulata</i> MURAY.					•			•				
146	<i>Holotrichia koraiensis</i> MURAY.					•							
* 147	<i>Eotrichia titanis</i> (REITT.)					•							
* 148	<i>Popillia indigonacea</i> MQTSCH.					•							
* 149	<i>Popillia atrocoerulea</i> BAT.	•	•			•			•				
150	<i>Popillia chinensis</i> FRIV.	•				•			•				
* 151	<i>Popillia japonica</i> NEWM.					•							
* 152	<i>Popillia quadriguttata</i> (FABR.)			•		•			•				
* 153	<i>Popillia ruficollis</i> KR.					•			•				
154	<i>Popillia uchidai</i> NIJ. et KIN.					•			•				
155	<i>Popillia quelpartiana</i> OH.					•			•				
* 156	<i>Phyllopertha horticola</i> (L.)					•							
157	<i>Phyllopertha diversa</i> WATERH.			•		•			•				•
* 158	<i>Spilota plagiocollis</i> FAIRM.					•			•				
* 159	<i>Proagopertha lucidula</i> (FALD.)					•			•				
* 160	<i>Mimela splendens</i> (GYLL.)			•		•			•				
* 161	<i>Mimela chinensis</i> KIRBY	•				•			•				

No	Species	A	B	C	D	E	F	G	H	I	K	L	Eurasia
162	<i>Mimela fusania</i> BAT.	•				•			•	•			
163	<i>Mimela concolor</i> BLANCH.	•	•			•				•	•		
164	<i>Mimela pekinensis</i> koreana MACH.					•			•				•
* 165	<i>Rhombonyx holosericea</i> (FABR.)			•	•	•	•	•	•				
166	<i>Rhombonyx testaceipes</i> MOTSCH.			•	•	•	•	•	•				
167	<i>Rhombonyx ussuriensis</i> MEDV.					•	•	•	•				
* 168	<i>Anomala (Chrysoplethisa) sieversi</i> HEYD.			•		•	•	•	•	•			
169	<i>Anomala (Chrysoplethisa) octiescostata</i> BURM.			•		•	•	•	•				
* 170	<i>Anomala (Euchronomala) cuprea</i> (HOPE)			•	•	•	•	•	•				
* 171	<i>Anomala (Euchronomala) viridana</i> KOLBE			•	•	•	•	•	•				
172	<i>Anomala (Euchronomala) mongolica</i> FALD.					•	•	•	•				
173	<i>Anomala (Euchronomala) albopilosa</i> HOPE		•			•	•	•	•		•		
174	<i>Anomala (Idiocnema) aulax</i> WIED.					•	•	•	•	•			
175	<i>Anomala (Idiocnema) costifera</i> REITT.					•	•	•	•	•	•		
176	<i>Anomala (Idiocnema) spiloptera</i> BURM.					•	•	•	•	•			
177	<i>Anomala (Anomala) pleurimargo</i> REITT.			•		•	•	•	•				
* 178	<i>Anomala (Anomala) corpulenta</i> MOTSCH.			•		•	•	•	•				
* 179	<i>Anomala (Anomala) luculenta</i> ER.					•	•	•	•				
180	<i>Anomala (Anomala) geniculata</i> MOTSCH.	•	•	•	•	•	•	•	•				
181	<i>Anomala (Anomala) rufocuprea</i> MOTSCH.				•	•	•	•	•				
182	<i>Anomala (Emphalena) chamaeleon</i> FAIRM.					•	•	•	•				
* 183	<i>Blitopertha (Exomala) pallidipennis</i> REITT.					•	•	•	•				
* 184	<i>Blitopertha (Exomala) orientalis</i> (WATERH.)		•	•	•	•	•	•	•				
* 185	<i>Blitopertha (Blitopertha) conspurcata</i> (HAR.)		•	•	•	•	•	•	•				
* 186	<i>Adoretus (Lepadoretus) sinicus</i> BURM.	•	•	•		•	•	•	•	•	•	•	
* 187	<i>Adoretus (Lepadoretus) tenuimaculatus</i> WATERH.	•	•	•		•	•	•	•	•	•	•	
* 188	<i>Rhomborrhina (Rhomborrhina) japonica</i> HOPE	•	•	•		•	•	•	•	•	•	•	
* 189	<i>Dicranocephalus adamsi</i> PASC.			•		•	•	•	•	•	•	•	
* 190	<i>Cetonia (Eucetonia) magnifica</i> BALL.					•	•	•	•				
191	<i>Cetonia (Eucetonia) viridiopaca</i> (MOTSCH.)				•	•	•	•	•				
192	<i>Cetonia (Eucetonia) pilifera</i> (MOTSCH.)			•	•	•	•	•	•				
193	<i>Cetonia (Eucetonia) roelofsi</i> HAR.			•	•	•	•	•	•				
* 194	<i>Polosia (Liocola) brevitarsis</i> (LEWIS)	•		•	•	•	•	•	•	•			
195	<i>Polosia (Liocola) lugubris orientalis</i> MEDV.			•	•	•	•	•	•				

No	Species	A	B	C	D	E	F	G	H	I	K	L	Eurasia
* 196	<i>Potosia (Calopotosia) aerata</i> (ER.)			•		•	•		•				
* 197	<i>Potosia (Potosia) fametica</i> (JANS.)				•	•	•	•	•				•
198	<i>Potosia (Potosia) metallica</i> (HBST)					•	•	•	•	•			
* 199	<i>Pocillophilides rusticola</i> (BURM.)			•		•	•	•	•				
* 200	<i>Glycyphana fulvitemma</i> (MOTSCH.)			•		•	•	•	•				
* 201	<i>Oryzctonia jucunda jucunda</i> (FALD.)	•		•		•	•	•	•				
* 202	<i>Clintrocera mandarina</i> (WESTW.)		•	•	•	•	•	•	•				
203	<i>Clintrocera obsoleta</i> (FAIRM.)					•	•	•	•				
* 204	<i>Dasyvalgus angusticollis</i> (WATERH.)		•	•	•	•	•	•	•				
205	<i>Altomyrina dichotomus</i> (L.)			•		•	•	•	•				
206	<i>Epiphileurus chinensis</i> (FALD.)		•	•	•	•	•	•	•			•	
207	<i>Oryctes rhinoceros</i> (L.)	•	•	•		•	•	•	•	•	•	•	
* 208	<i>Lasiotrichius succinctus</i> (PALL.)		•	•	•	•	•	•	•			•	
* 209	<i>Gnorimus subopacus</i> MOTSCH.		•	•	•	•	•	•	•			•	
210	<i>Osmoderma barnabita</i> MOTSCH.				•	•	•	•	•				
* 211	<i>Trox sabulosus</i> OLIV.					•	•	•	•				•
212	<i>Trox scaber</i> (L.)		•	•	•	•	•	•	•				•
213	<i>Trox opacotuberculatus</i> MOTSCH.			•		•	•	•	•				•
214	<i>Nigidius miwai</i> NAG.			•		•	•	•	•				
215	<i>Prismagnathus dauricus</i> MOTSCH.					•	•	•	•	•			
* 216	<i>Psilidoremus inclinator</i> (MOTSCH.)					•	•	•	•				
* 217	<i>Macrodercas rubrofemoratus</i> (VOLL.)	•	•	•	•	•	•	•	•				
218	<i>Macrodercas rectus</i> (M TSCH.)	•	•	•	•	•	•	•	•				
* 219	<i>Macrodercas stritipennis</i> MOTSCH.	•	•	•	•	•	•	•	•				
* 220	<i>Serrogathus titanus</i> (BOISD.)	•	•	•	•	•	•	•	•				
* 221	<i>Dorcus hopei</i> (SAUND.)	•	•	•		•	•	•	•	•	•	•	
* 222	<i>Dorcus montivagus</i> (LEVIS)	•	•	•		•	•	•	•	•	•	•	
223	<i>Lucanus maculifemoratus</i> MOTSCH.		•	•	•	•	•	•	•				
224	<i>Eurytrachelidius haitichunus</i> DID. et SEG.	•	•	•	•	•	•	•	•				


ern species succeeding into south in the cold periods. Local fauna has been formed gradually; it was composed of various groups which found there favorable conditions for survival. In the core of the Far-Eastern forest fauna of *Scarabaeoidea* dominating in the Korean Peninsula, occur some further changes. The elements that were previously widely distributed, nowadays occur locally in the mainstay areas of the Tertiary flora. Many of the phytophagous species have adapted themselves to the present conditions of the plant landscape created by man. They follow the succession of the agriculture and undergo gradual synantropization. The coprophagic species, majority of which is peculiar to the open areas, occupy large territories, in compliance with physical properties of the soil and their own adaptative abilities. Penetration of the Peninsula by the ubiquitous species, elements of North-Chinese, Manchurian, South-Siberian and Mongolian fauna as well as by Oriental species, will possibly develop. The main, most convenient pathways of migration are in the coastal regions; they lead from China and Manchuria at the Gulf of Korea and from Primorskij Kraj along the Japanese Sea.

The summing-up of the above conclusions indicate that the fauna of the Korean Peninsula is considerably complex; its elements join the Far-Eastern faunas of *Scarabaeoidea*, which have common history and origin and occur in the similar environmental conditions. The separateness of this fauna from those of the rest of Palaearctic is evident. Nevertheless, the determination of the zoogeographic border between both Far-Eastern, Palaearctic and Oriental faunas is not possible. The transitory zone of both types is very wide; their overlap is various in different regions of Far East. Consequently, with appropriateness to the distribution of various animal groups, their ecologic requirements and abilities to pass the barriers, the transition is formed somewhat in a different way.

LIST OF THE KOREAN GEOGRAPHIC PLACE-NAMES MENTIONED IN THE TEXT

Numbers indicate the localities on the map, Fig. 3. Japanese transcription of some place-names is given in square brackets.

1 — Anak [Angaku]	38°31', 125°28'
2 — Andžu [Anshu]	39°35', 125°38'
3 — Čedžu-do [Saishu-to, Saisyu-to] (Quelpart Island). A large island in the East China Sea off the southern tip of Korea	
4 — Čungdžu [Chuschu]	36°58', 127°57'
5 — Čočhingvŏn [Chochiin]	36°36', 127°21'
6 — Čongdžu [Seisyu, Seishu]	35°35', 126°50'
7 — Čhŏngan [Tenan]	36°49', 127°11'
8 — Čhunčhŏn [Shunsen, Syunsen]	37°52', 127°44'
9 — Čongsŏng [Shojo]	37°24', 128°39'
10 — Čongphjŏng	37°09', 129°14'
11 — Čhŏngdžin [Seishin, Seisin]	41°47', 129°48'
12 — Čhŏngdžu [Zeishu, Zensyu]	36°38', 127°38'



13 — Chonma-san, Pakjŏn	near: 37°51', 126°35'
14 — Čhŏnne	39°28', 127°13'
15 — Čojang	near: 39°50', 127°30'
16 — Čangdžin-ho [Cho-shin-ko] Nangnim-sanmek	40°25', 127°09'
17 — Džuyr Jonghen	near: 41°40', 129°35'
18 — Hedžu [Kaishu, Kaisyu] Hakhjŏn-čhŏn	38°03', 125°46'
19 — Hamhyng [Kanko] Tŏksan-ri	39°55', 127°34'
20 — Hongčhŏn [Kosen]	37°41', 127°54'
21 — Hadong [Kato]	35°05', 127°44'
22 — Hvangjang	38°43', 127°36'
23 — Hvačhŏn	39°02', 126°05'
24 — Henam [Kainan]	34°35', 126°40'
25 — Hverjŏng [Kaiinji]	42°27', 129°41'
26 — Hjesan [Keisanchin, Kaizanchin] Dŏngha-ri, Džedang-jong	41°23', 128°13'
27 — Hjangsan Mjohjang-san, Manphok-tong, Munsu-tong, Hapiro, Hjangam-ri, Hjangsan-čhŏn	40°01', 126°12'
28 — Hyngnam	39°50', 127°39'
29 — Hamdžu Hyeongpong-ri	near: 39°45', 127°30'
30 — Hongvŏn [Kogen] Jŏnpong-ri	40°04', 127°56'
31 — Hyngsang	39°53', 127°26'
32 — Inčhŏn (Chemulpo) [Jinsen, Zinsen]	37°28', 126°40'
33 — Ičhŏn [Risen]	38°28', 126°52'
34 — Jongdžu [Eishu]	36°50', 128°37'
35 — Jončhŏn [Eisen]	38°06', 127°06'
36 — Jongdŏk [Eido]	36°24', 129°24'
37 — Jongvŏn [Neietsu]	39°49', 126°34'
38 — Jŏsu [Reisui]	34°45', 127°45'
39 — Jodžu [Reishu]	37°18', 127°40'
40 — Jongak-san Sunha-gang	near: 38°50', 125°20'
41 — Jongsŏng Maram	39°10', 125°48'
42 — Kangnyng [Koryo, Korjo]	37°45', 128°53'
43 — Kvangjang	34°58', 127°36'
44 — Kjŏngsŏng [Kyojo] Poro-čhŏn, Mehjang-ri	41°34', 129°38'
45 — Kangge [Kokai]	40°57', 126°37'
46 — Kunsan [Gunsan, Gunzan]	35°58', 126°42'
47 — Kvangdžu [Koshu, Kosyu]	35°09', 126°53'
48 — Kočhang [Kyosho]	35°26', 126°38'
49 — Kangjasan [Kyokosan]	35°47', 127°06'
50 — Kangdžin [Koshin]	34°38', 126°44'

Fig. 3. Schematic map of the Korean Peninsula. Numbers mark the localities specified in the list

51 — Kjongdžu [Keishu, Keisyu]	35°49', 129°14'
52 — Kimpho	37°37', 126°41'
53 — Kymgang-san Masin-rjǒng	38°37', 128°07'
54 — Kosan [Kosen]	35°57', 127°16'
55 — Kildžu [Kisshu, Kissyu]	40°57', 129°21'
56 — Kesǒng [Kaijo]	37°56', 126°33'
57 — Kečhǒn	39°43', 125°53'
58 — Kymčhǒn	38°09', 126°27'
59 — Kodže [Kotei]	34°51', 128°36'
60 — Kvanmo-bong [Kwanboho]	41°42', 129°15'
61 — Kangsǒ [Kosi, Kosei] Thesǒng	38°58', 125°29'
62 — Kvesan [Kwasan]	36°49', 127°47'
63 — Komusan [Komosan]	42°07', 129°43'
64 — Koksan [Kokuzan]	38°46', 126°39'
65 — Kudžang Džosan-ri	39°52', 126°04'
66 — Mokpho [Moppo]	34°48', 126°24'
67 — Mudžu [Moshu, Mosyu]	36°01', 127°41'
68 — Mjǒngčhǒn [Meisen]	41°04', 129°28'
69 — Mengsan	39°40', 126°29'
70 — Musan [Mosan]	42°12', 129°14'
71 — Munčhǒn [Bunsen]	39°14', 127°23'
72 — Mačǒn	near: 39°55', 127°50'
73 — Namvǒn [Nangen]	35°24', 127°22'
74 — Nadžin [Rashin, Rasin]	42°14', 130°15'
75 — Nampho [Chinnampo, Tinnampo] Tedong-gang, Usan-ri, Vaudo	38°45', 125°25'
76 — Namphote-san	41°44', 128°24'
77 — Onpho-ri	near: 41°32', 129°25'
78 — Oro	40°02', 127°29'
79 — Pujǒ [Fuyo, Huyo]	36°18', 126°54'
80 — Pektu-san [Hakuyosan, Hakutosan]	41°56', 128°08'
81 — Phohang [Hoko]	36°03', 129°20'
82 — Pegam (Not localized on the map)	
83 — Pusan [Fusan, Husan]	35°09', 129°02'
84 — Peksan	39°30', 126°42'
85 — Pungan [Hozan]	35°43', 126°43'
86 — Phjǒngjang [Heijo] Tedong-gang, Mankjǒngde, Mankjǒng-bong, Moran-bong	39°00', 125°45'
87 — Purjǒng Thomak-tong, Musu-ri, Hjǒngčǎe-tong	42°05', 129°43'
88 — Ponghva-ri	near: 39°00', 125°25'
89 — Pisylsan (Not localized on the map)	
90 — Počhǒn Počhǒnbo, Karim-čhǒn, Posǒ-ri	41°32', 128°20'
91 — Pučhǒn-gang [Fusen-ko]	near: 40°40', 127°30'
92 — Suvǒn [Suigen]	37°17', 127°01'
93 — Soul [Keijo, Keizyo]	37°32', 127°00'
94 — Sunčhǒn [Juntan] Džamo-ri, Džamo-san	near: 39°25', 125°40'

95 — Sŏnchŏn [Seisen]	39°46', 124°51'
96 — Sinyidžu [Shingishu, Singisyu]	40°06', 124°23'
97 — Sičhang	40°08', 128°28'
98 — Sepho	38°39', 127°26'
99 — Sinphjŏng	38°53', 126°44'
100 — Samchŏnpho [Sanseizan]	34°56', 128°05'
101 — Sarivŏn [Sharei, Shariin]	38°30', 125°48'
102 — Sungho [Shoko]	near: 38°17', 125°22'
103 — Samdžijŏn	41°52', 128°23'
Phote-čhŏn, Posŏ-ri	
104 — Sunan	39°14', 125°40'
Junha-ri	
105 — Sŏkam-Čösudži	near: 39°20', 125°42'
106 — Samsŏk	near: 39°10', 125°40'
Tečhŏn-ri, Sŏngmun-ri	
107 — Sičung-ho	near: 39°05', 127°24'
108 — Sinčhŏn [Shinsen]	38°20', 125°30'
109 — Sangvŏn-am	near: 39°55', 126°20'
110 — Susŏng-čhŏn	near: 41°54', 129°43'
111 — Tongne [Torai]	35°12', 129°06'
112 — Tegu [Taikyu]	35°52', 128°35'
113 — Tedžon [Taisei]	36°18', 127°26'
114 — Thosan	38°19', 126°42'
115 — Tongčong-ho	near: 39°00', 127°30'
116 — Usi	40°40', 125°40'
117 — Uldžin	37°00', 129°24'
118 — Vŏnsan [Gensan, Genzan]	39°10', 127°28'
Ungč'in	
119 — Vondžu [Genshu]	37°20', 127°59'
120 — Vando	34°19', 126°43'
121 — Sujang-san	38°06', 125°40'
122 — Kosŏng [Kojo]	38°45', 128°12'
123 — Sam-il-pho	near: 38°40', 128°12'
124 — Tesŏng-san	near: 39°06', 125°52'
125 — Chosan [Sozan]	40°50', 125°46'
126 — Korjŏng	35°44', 128°17'
127 — Hvado	33°42', 126°23'
128 — Sinsang	39°40', 127°27'
129 — Sokčhŏ	38°15', 128°38'
130 — Sŏnsan	36°15', 128°09'
131 — Kongdžu [Koshu]	36°25', 127°08'
132 — Nungni	38°40', 126°11'

Scarabaeinae

Scarabaeini

Scarabaeus (Scarabaeus) affinis BRULLÉ, 1832

Distribution. South Europe, Asia Minor, Central Asia, Afghanistan, Tibet, West and North China, Manchuria, Korea (BALTHASAR, 1963 et all.).

Material examined. 1♂. Prov. Hvanghe-namdo, Hedžu (18), 1 VI 1974, at light, Z. STEBNICKA (ISEZ).

Remarks. The specimen collected is a representative of var. *typhon* FISCH., which occurs in the eastern part of the distributional area of this species.

Gymnopleurini

Gymnopleurus (Gymnopleurus) mopsus (PALLAS, 1781)

Distribution. South Europe, North Africa, Transcaucasus, Asia Minor Central Asia, Mongolia, North and Central China (BALTHASAR, 1963; ENDRÖDI 1971 et all.).

Material examined. 21 ♂♂ and ♀♀. Prov. Hamgjöng-namdo, distr. Hongvön (30), Jönpong-ri, 8 VI 1965, in cow dung, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Hamhyng-si, distr. Hamdžu (29), Hyngpong-ri, 12 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. Palaearctic species, new to the Korean fauna.

Gymnopleurus (Paragymnopleurus) sinuatus (OLIVIER, 1789)

(Fig. 4)

Distribution. Central and South China, Taiwan (Rosen, Nanshanchi, Wushe, Taloko, Lutao Is., Lanyu Is.), India, Indochina (Thailand), Sunda Islands (BALTHASAR, 1963; NOMURA, 1973).

Material examined. 1 ♀. Prov. Hamgjöng-namdo, distr. Hongvön (30), Jönpong-ri, 8 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. Species of the Oriental origin, new to the Korean fauna.

Sisyphini

Sisyphus (Sisyphus) schaefferi (LINNAEUS, 1758)

Distribution. Central and South Europe, North Africa, Asia Minor, Central Asia, Mongolia, North China, Korea (BALTHASAR, 1963 et all.).

Material examined. 6 ♂♂ and ♀♀. Prov. Hamgjöng-namdo, distr. Hongvön (30), Jönpong-ri, 8 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Coprinae

Oniticellini

Liatongus (Liatongus) phanaeoides (WESTWOOD, 1840)

(Fig. 5)

Distribution. China, Taiwan (Nanshanchi, Funchiifo), India, Indochina, Korea, Japan — Hokkaido, Honshu, Sado (Oda, Kitauzima, Mt. Kinpoku, Aoneba-goe, Ookura-goe), Shikoku, Kyushu, Ryu-kyu Isl. (NAKANE et BABA, 1960; NAKANE, 1961; BALTHASAR, 1963; NOMURA, 1973).

Material examined. 56 ♂♂ and ♀♀. Japan — Honshu (Chiba, Owada), 8 VII 1937 (HMNH); Korea — Prov. Hamgjong-namdo, distr. Hjangsan (27), Mjohjang-san (Mts), 16 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), distr. Hongvön (30), Jönpöng-ri, 8 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Hamgjong-pukto, distr. Kjongsöng (44), Onpho-ri (77), mountain valley, 21 VIII 1959, B. PISARSKI and J. PRÓSZYŃSKI (ZIW), vicinity of Džuyr (17), dry sunny hill, 5 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW); Prov. Kangvön-do, Kymgang-san (Mts, 53), 16—18 VI 1974, in cow dung, Z. STEBNICKA (ISEZ), 34 km W of Vönsan (118), mountain pass Masin-rjöng, 16 IX 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. This species occurs exclusively in the mountainous regions and in mountains where it reaches 2500 m above sea level.

Copris (Copris) ochus (MOTSCHULSKY, 1860)

Distribution. North- and Central China, Mongolia, Manchuria, Korea (Čedžu-do, 3), Japan — Miyake, Hokkaido, Honshu, Sado (Sawada), Shikoku, Kyushu, Iki (NAKANE et BABA, 1960; BALTHASAR, 1963; NOMURA, 1969).

Material examined. 12 ♂♂ and ♀♀. Prov. Phjongan-namdo, distr. Kangsö (61), Thesöng, vicinity of the lake, 24 VIII 1971, collected at light of UV lamps, J. RAZOWSKI (ISEZ); Prov. Phjongsjang-si, Ponghva-ri (88), 36 km W of Phjongsjang (86), 24 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), Jongak-san (Mt. 40), 20 VIII 1971, collected at light of UV lamps, J. RAZOWSKI (ISEZ); Prov. Hvanghe-namdo, Sinčhön (108), valley of river, 16 IX 1971, in cow dung, J. PAWŁOWSKI (ISEZ), Hedžu (18), 3 VI 1974, at light of UV lamps, Z. STEBNICKA (ISEZ).

Copris (Copris) tripartitus WATERHOUSE, 1875

Distribution. China, Korea (Čedžu-do, 3), Taiwan (Nanshanchi), Japan — Honshu, Kyushu, Tsushima (BALTHASAR, 1963; NOMURA, 1973).

Material examined. 11 ♂♂ and ♀♀. Prov. Phjongan-namdo, distr. Kangsö (61), vicinity of the lake, 13 VI 1974, in cow dung, Z. STEBNICKA (ISEZ), 19 IX 1966, H. SZELEGIEWICZ and C. DZIADOSZ (ZIW), 24 VIII 1971, at light of UV lamps, J. RAZOWSKI (ISEZ), distr. Sunčhön (94), Džamo-san, 27 VIII 1971, at light of UV lamps, J. PAWŁOWSKI (ISEZ); Prov. Kesöng-si, vicinity of the water-fall Pakjön on the Čhönma-san (Mt. 13), 5 VI 1974, in cow dung, Z. STEBNICKA (ISEZ).

Caccobius (Caccophilus) christophi HAROLD, 1879

(Fig. 6)

Distribution. USSR — Amur, Ussuri, Primorskij Kraj; North and Central China, Korea (BALTHASAR, 1963).

Material examined. 30 ♂♂ and ♀♀. Prov. Hamgjong-pukto, distr. Kjongsöng (44), Onpho-ri (77), 20 IX 1959, B. PISARSKI and J. PRÓSZYŃSKI (ZIW);

Prov. Janggang-do, distr. Hjesan (26), Dŏngha-ri, 2 IX 1971, in cow dung, J. PAWŁOWSKI (ISEZ); Prov. Hamgjöng-namdo, distr. Hjangsan (27), valley Hapiro in the Mjohjang-san (Mts), 20—21 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. Manchurian species, local and not frequent in D.P.R.K.

***Caccobius (Caccophilus) sibiricus* BALTHASAR, 1935**

(Fig. 7)

Distribution. USSR — Transbaikal Region, Ussuri, Primorskij Kraj (Vladivostok) (BALTHASAR, 1963).

Material examined. 26 ♂♂ and ♀♀. Prov. Janggang-do, distr. Samdžijon (103), valley of river Phote-čhŏn, 8 IX 1971, in cow dung, J. PAWŁOWSKI (ISEZ), distr. Hjesan (26), Dŏngha-ri, 2 IX 1971, in cow dung, A. SZEPTYCKI (ISEZ).

Remarks. Species new to the Korean fauna, probably local. Single population found in one of the northern Provinces. Occurs in cattle dung.

***Caccobius (Caccophilus) sordidus* HAROLD, 1886**

(Fig. 10)

Distribution. USSR — Ussuri, Primorskij Kraj; Manchuria (Harbin), North China, Korea (BALTHASAR, 1963).

Material examined. 36 ♂♂ and ♀♀. Prov. Janggang-do, distr. Hjesan (26), Dŏngha-ri, 2 IX 1971, in cow dung, J. PAWŁOWSKI (ISEZ); Prov. Čhŏngdžin-si, valley of river Susŏng-čhŏn (110), 21 V 1974, Z. STEBNICKA (ISEZ), distr. Purjŏng (87), 22 V 1974, J. PAWŁOWSKI (ISEZ).

***Caccobius (Caccophilus) brevis* WATERHOUSE, 1875**

(Fig. 9)

Distribution. East Siberia (Ussuri), North China, Manchuria, Korea (Čedžu-do, 3) (BALTHASAR, 1963).

Material examined. 89 ♂♂ and ♀♀. Prov. Hamgjöng-punkto, distr. Kjöngsŏng (44), Onpho-ri (77), 20 VIII 1959, B. PISARSKI and J. PRÓSZYŃSKI (ZIW); Prov. Phjŏngjang-si, Ponghva-ri (88), 24 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW); Prov. Janggang-do, distr. Hjesan (26), Dŏngha-ri, 2 IX 1971, J. PAWŁOWSKI (ISEZ); Prov. Čhŏngdžin-si, valley of river Susŏng-čhŏn (110), 2 IX 1971, J. PAWŁOWSKI (ISEZ); Prov. Kesŏng-si, vicinity of the water-fall Pakjŏn on the Čhŏnma-san (Mt. 13), 5 VI 1974, in cow dung and at human excrements, Z. STEBNICKA (ISEZ).




Fig. 4. *Gymnopleurus* (*Paragymnopleurus*) *sinuatus* (OLIV.) — female head. Figs. 5—12. Male genitalia, lateral and dorsal view; 5 — *Liatongus* (*Liatongus*) *phanaeoides* (WESTW.); 6 — *Caccobius* (*Caccophilus*) *christophi* HAR.; 7 — *C. (C.) sibiricus* BALTH.; 8 — *C. (C.) unicornis* (FABR.); 9 — *C. (C.) brevis* WATERH.; 10 — *C. (C.) sordidus* HAR.; 11, 12 — *Onthophagus* (*Digitonthophagus*) *solivagus* HAR.

***Caccobius* (*Caccophilus*) *unicornis* (FABRICIUS, 1798)**

Syn.: *Caccobius yamauchii* MATSUMURA, 1936.

(Fig. 8)

Distribution, Japan — Honshu, Kyushu, Ryu-kyu Isl.; Korea, China, Taiwan (Liyutan, Roshan, Funchiifo, Chipon, Oulanpi), India, Burma, Indochina, Malaysia, Sunda Isl., Philippines (BALTHASAR, 1963; NOMURA, 1973).

Material examined. 11 ♂♂ and ♀♀. Prov. Hvanghe-namdo, valley of river

Hakhjön-čhön near Hedžu (18), 4 VI 1974, in sheep dung, Z. STEBNICKA (ISEZ); Prov. Kangvön-do, Kymgang-san (Mts, 53), 16 VI 1974, at human excrements, Z. STEBNICKA (ISEZ).

***Onthophagus (Digitonthophagus) solivagus* HAROLD, 1886**

(Figs. 11, 12)

Distribution. North- and Central China, Korea, (BALTHASAR, 1963).

Material examined. 58 ♂♂ and ♀♀. Prov. Hamgjöng-punkto, distr. Kjön-gsöng (44). Onpho-ri (77), 20 VIII 1959, B. PISARSKI and J. PRÓSZYŃSKI (ZIW); Prov. Hamgjöng-namdo, distr. Hjangsan (27), valley Munsu-tong in the Mjohjang-san (Mts), 18 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), distr. Hongvön (30), Jönpong-ri, 8 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Phjongan-namdo, distr. Sunan (104), Sökam-Čösudži (105), 30 V 1974, in sheep dung, Z. STEBNICKA (ISEZ), Junha-ri, 13 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW); Prov. Phjôngjang-si, Phjôngjang (86), 16 V 1974, at human excrements, Z. STEBNICKA (ISEZ); Prov. Čhöngdžin-si, valley of river Susöng-čhön (110), 25 V 1974, in cow dung, Z. STEBNICKA (ISEZ); Prov. Kesöng-si, Čhönma-san (Mt., 13), 6 VI 1974, Z. STEBNICKA (ISEZ); Prov. Hvanghe-namdo, vicinity of the water-fall Sujang-san (121), 2 VI 1974, at human excrements, Z. STEBNICKA (ISEZ); Prov. Kangvön-do, Vönsan (118), 15—20 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), Kymgang-san (Mts, 53), 17 VI 1974, Z. STEBNICKA (ISEZ).

Remarks. This species is probably an element of Manchurian fauna; it has relatively narrow area of distribution. Frequent, in some Provinces numerous and widely distributed.

***Onthophagus (Digitonthophagus) hvangeus* STEBNICKA, 1973**

(Fig. 13)

Distribution. North Korea — Prov. Hvanghe-namdo, Sinčhön (108) (STEBNICKA, 1973).

Material examined. 6 ♂♂ and ♀♀. Prov. Phjôngjang-si, Jongak-san (Mt., 40), valley of river, 29 V 1974, in sheep dung, Z. STEBNICKA (ISEZ).

Remarks. Only one male specimen of this species has been known up to the present. The female differs from the male by the shape of anterior tibia and by strongest and more distinct clypeal punctures, which are transversely wrinkled near anterior margin.

***Onthophagus (Strandius) lenzi* HAROLD, 1874**

(Figs. 14, 15)

Distribution. North- and Central China, Manchuria, Korea, Taiwan, Japan — Hokkaido, Honshu, Sado, Shikoku, Kyushu, Tsushima, Nakanoshima, Tanegashima, Ohshima, Toshima, Miyake, Mikura, Hachijo, Hachijo-Kojima, Aogashima (BALTHASAR, 1963; NOMURA, 1966; 1969).

Material examined. 160 ♂♂ and ♀♀. Prov. Phjǽngan-namdo, distr. Sunan (104), Junha-ri, 40 km W of Phjǽngjang (86), 13 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), Sǽkam-Čǽsudŷi (105), 27 VIII 1971, on the banks of lake, at light of UV lamps, J. PAWŁOWSKI (ISEZ), 30 V 1974, in cow dung, Z. STEBNICKA (ISEZ), distr. Kangsǽ (61), Thesǽng, 26 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), 24 VIII 1971, at light of UV lamps, J. RAZOWSKI (ISEZ), distr. Sunčǽn (94), Dŷamo-san, 27 VIII 1971, at light of UV lamps, J. PAWŁOWSKI (ISEZ); Prov. Phjǽngjang-si, distr. Jongsǽng (41), Maram, 20 km NE of Phjǽngjang, 29 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), Ponghva-ri (88), 36 km E of Phjǽngjang, 24 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), Mankjǽngde, 31 VIII 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), Phjǽngjang (86), 27 VIII 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), 17 V 1974, at light of UV lamps, Z. STEBNICKA (ISEZ), Moran-bong (municipal park), at human excrements, J. PAWŁOWSKI and A. SZEPTYCKI (ISEZ), Botanical Garden, 15 IX 1971, J. PAWŁOWSKI (ISEZ), Jongak-san (Mt., 40), 20 VIII 1971, at light of UV lamps, J. PAWŁOWSKI (ISEZ), 17 VIII 1971, J. RAZOWSKI (ISEZ), 29 V 1974, in sheep dung, Z. STEBNICKA (ISEZ); Prov. Kangvǽn-do, Vǽnsan (118), 1 IX 1966, H. SZEŁĘGIEWICZ and C. DZIADOSZ (ZIW), 15—20 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), Kymgang-san (Mts, 53) near Kosǽng (122), 16 VI 1974, in cow dung, Z. STEBNICKA (ISEZ); Prov. Hvanghe-namdo, Sinčǽn (108), 16 IX 1971, in cow dung on the banks of river, J. PAWŁOWSKI (ISEZ), vicinity of the water-fall Sujang-san (121) near Hedŷu (18), 2 VI 1974, at human excrements, Z. STEBNICKA (ISEZ), valley of river Hakhjǽn-čǽn near Hedŷu (18), 3 VI 1974, in cow dung, Z. STEBNICKA (ISEZ).

Remarks. Very common species in D.P.R.K., locally frequent, occurs in various biotopes. Found under excrements in the soil up to 10 cm below its surface. Readily comes to light; numerous specimens have been collected under the street lamps.

Onthophagus (Strandius) japonicus HAROLD, 1874


(Figs. 18, 19)

Distribution. Korea, Japan — Honshu, Sado (Oda), Taiwan (Wushe) (NAKANE et BABA, 1960; BALIHASAR, 1963; NOMURA, 1973).

Material examined. 28 ♂♂ and ♀♀. Prov. Hamgjǽng-pukto, distr. Kjǽngsǽng (44), valley Onpho-ri (77), 3 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), Jonghen near Dŷuyr (17), 25 V 1974, in cow dung, Z. STEBNICKA (ISEZ); Prov. Hamgjǽng-namdo, distr. Hongvǽn (30), Jǽnpong-ri, 8 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Phjǽngjang-si, distr. Samsǽk (106), Sǽngmun-ri, 22 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), Poghva-ri (88) 36 km W of Phjǽngjang, 24 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW); Prov. Phjǽngan-namdo, distr. Sunan (104), Junha-ri 40 km W of Phjǽn-

gjang, 13 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), Sökam-Čösu-dži (105), 30 V 1974, in cow dung on the banks of lake, Z. STEBNICKA (ISEZ); Prov. Kesöng-si, Čhönma-san (Mt., 13), 8 VI 1974, in cow dung, Z. STEBNICKA (ISEZ); Prov. Čhöngdžin-si, valley of river Susöng-čhön (110), 21 V 1974, in cow dung, Z. STEBNICKA (ISEZ); Prov. Hvanghe-namdo, valley of river Hakhjön-čhön near Hedžu (18), 3 VI 1974, in cow dung, Z. STEBNICKA (ISEZ), Sujang-san (Mt., 121), 3 VI 1974, at human excrements, Z. STEBNICKA (ISEZ).

Remarks. According to BALTHASAR (1963) this species is not frequent but widely distributed in Japan; its similar distribution has been observed in D.P.R.K., where single specimens often occur in various biotopes.



Figs. 13—21. Male genitalia, lateral and dorsal view: 13 — *Onthophagus (Digitonthophagus) hvangheus* STEB.; 14, 15 — *O. (Strandius) lenzi* HAR.; 16, 17 — *O. (Phanaeomorphus) fodiens* WATERH.; 18, 19 — *O. (Strandius) japonicus* HAR.; 20, 21 — *O. (Gibbonthophagus) viduus* HAR. Fig. 22. *O. (Phanaeomorphus) ater* WATERH. — female head

Onthophagus (Phanaeomorphus) fodiens WATERHOUSE, 1875

(Figs. 16, 17)

Distribution. North- and Central China, Manchuria, Korea, Japan — Honshu (Kurokawa, Takada, Mt. Ninooji, Matsunoyama, Nagaoka-Hachibu-se), Sado (Umezū, Futami, Ogura-toge) (NAKANE et BABA, 1960; BALTHASAR, 1963).

Material examined. 30 ♂♂ and ♀♀. Prov. Hamgjong-namdo, distr. Hongvŏn (30), Jŏnpong-ri, 8 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Kangvŏn-do, Kymgang-san (Mts, 53) near Kosŏng (122), 16—18 VI 1974, at human excrements, Z. STEBNICKA (ISEZ); Prov. Čhŏngdžin-si, valley of river Susŏng-čhŏn (110), 21 V 1974, in cow dung, Z. STEBNICKA (ISEZ); Prov. Hvanghe-namdo, Sujang-san (Mt., 121), 2 VI 1974, at human excrements, Z. STEBNICKA (ISEZ), valley of river Hakhjŏn-čhŏn near Hedžu (18), 3 VI 1974, in cow dung, Z. STEBNICKA (ISEZ); Prov. Kesŏng-si, vicinity of the water-fall Pakjŏn on the Čhŏnma-san (Mt., 13), 6 VI 1974, in cow dung, Z. STEBNICKA (ISEZ); Prov. Phjŏngan-namdo, distr. Nampho (75), locality Vaudo at the estuary of the river Tedong-gang, 18 IX 1971, under stone, J. PAWŁOWSKI (ISEZ), distr. Sunan (104), Sŏkam-Čŏsudži (105), 21 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Phjŏngjang-si, Tesŏng-san (Mts, 124), 18 V 1974, in cow dung, Z. STEBNICKA (ISEZ); Prov. Hamhyng-si, Hamhyng (19), 13 IX 1966, H. SZELEGIEWICZ and C. DZIADOSZ (ZIW).

Remarks. Species widely distributed but not frequent, found as single specimens in various biotopes. According to the NOMURA's data (1973), the other similar species named *O. yubarinus* MATS. (= *Matasha mushana* MATS. sensu NOMURA, 1973) probably occurs in Taiwan; BALTHASAR (1963) considered it as a synonym of *O. fodiens* WATERH.

Onthophagus (Phanaeomorphus) ater WATERHOUSE, 1875

(Fig. 22)

Distribution. East Siberia, North- and Central China (Sichuan), Korea, Taiwan (Sungkang, Funchiifo), Japan — Hokkaido, Honshu (Kurokawa, Matsunoyama, Midama), Awa-shima, Sado (Oda, Hatano, Kitauzima, Mt. Donden, Ogura-toge), Shikoku, Kyushu, Yakushima, Miyake, Mikura, Hachijo (NAKANE et BABA, 1960; BALTHASAR, 1963; NOMURA, 1969; 1973).

Material examined. 1♀. Prov. Hvanghe-namdo, vicinity of the water-fall Sujang-san (211), 2 VI 1974, at human excrements, Z. STEBNICKA (ISEZ).

Onthophagus (Gibbonthophagus) viduus HAROLD, 1874

(Figs. 20, 21)

Distribution. North China (Shandong), Manchuria, Korea, Japan — Hokkaido, Honshu (Kurokawa, Niigata, Sakasamaki), Sado (Nakaoku), Shikoku, Kyushu, Yakushima, Tanegashima, Tsushima, Nakanoshima, Takarajima, Ama-

mi-Ōshima, Okinawa, Ishigaki, Iriomote, Ohshima, Niijima, Miyake, Hachijo (NAKANE et BABA, 1960; BALTHASAR, 1963; NOMURA, 1966; 1969).

Material examined. 1 ♂ and 2 ♀♀. Prov. Hvanghe-namdo, vicinity of the water-fall Sujang-san (121), 2 VI 1974, at human excrements, Z. STEBNICKA (ISEZ).

***Onthophagus (Gibbonthophagus) atripennis* WATERHOUSE, 1875**

(Figs. 23, 24)

Distribution. North- and Central China (Sichuan), Manchuria, Korea, Japan — Honshu (Kurokawa, Hanadate), Sado (Iwakubi, Ookura-goe, Akado mari, Aono-toge), Shikoku, Kyushu, Tsushima, Tanegashima, Mikura (NAKANE et BABA, 1960; BALTHASAR, 1963; NOMURA, 1969).

Material examined. 5 ♂♂ and ♀♀. Prov. Hvanghe-pukto, Sarivŏn (101), 20—28 VIII 1966, M. MAGYAR (HMNH); Prov. Phjŏngjang-si, Phjŏngjang (86), 1 VII 1950, BORCHSENIUS (ZIL); Prov. Hamgjŏng-namdo, Nangnim-san-mek (Mts, 16), 7 VIII 1950, BORCHSENIUS (ZIL).

***Onthophagus (Onthophagus) uniformis* HEYDEN, 1886**

(Figs. 25, 26)

Distribution. East Siberia (Amur, Ussuri), Manchuria (BALTHASAR, 1963).

Material examined. 6 ♂♂ and ♀♀. USSR — Primorskij Kraj (Vladivostok) (HMNH); Korea — Prov. Hamgjŏng-namdo, distr. Hjangsan (27), valley Hapiro in the Mjohjang-san (Mts), 20—21 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Janggang-do, distr. Hjesan (26), Džedang-jong, 31 VIII 1971, at light of UV lamps, J. RAZOWSKI (ISEZ).

Remarks. Manchurian species of probably small area of distribution. New to the Korean fauna.


***Onthophagus (Onthophagus) gibbulus* (PALLAS, 1781)**

(Fig. 27)

Distribution. South- and Central Europe, Transcaucasus, Asia Minor, Central Asia, Mongolia, Siberia (Bajkal Lake, Amur, Ussuri), Manchuria, North China (BALTHASAR, 1963; ENDRÖDI 1967; 1968; 1969; 1971; TESÁŘ, 1968; MEDVEDEV, 1976).

Material examined. 32 ♂♂ and ♀♀. Prov. Janggang-do, distr. Hjesan (26), valley Dongha-ri, 2 IX 1971, in cow dung, J. PAWŁOWSKI (ISEZ); Prov. Hamgjŏng-pukto, distr. Kjŏngsŏng (44), valley Onpho-ri (77), 10 VIII 1959, B. PI-SARSKI and J. PRÓSZYŃSKI (ZIW), 7 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW); Prov. Phjŏngan-namdo, distr. Sunan (104), Junha-ri 40 km W of Phjŏngjang (86), 13 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW);

Prov. Phjǽngjang-si, Ponghva-ri (88), 24 IX 1970, R. BIELAWSKI and M. MROCKOWSKI (ZIW); Prov. Čhǽngdžin-si, distr. Purjǽng (87), 22 V 1974, in sheep dung on the banks of river, Z. STEBNICKA (ISEZ); Prov. Hvanghe-namdo,



Figs. 23—34. Male genitalia, lateral and dorsal view; 23, 24 — *Onthophagus (Gibbonthophagus) atripennis* WATERH.; 25, 26 — *O. (Onthophagus) uniformis* HEYD.; 27 — *O. (O.) gibbulus* (PALL.); 28, 29 — *O. (O.) bivertex* HEYD.; 30, 31 — *O. (O.) olsoufieffi* BOUC.; 32 — *O. (O.) punctator* REITT.; 33, 34 — *O. (O.) rugulosus* HAR.

Sinčhŏn (108), valley of river, 16 IX 1971, in cow dung, A. SZEPTYCKI (ISEZ).

Remarks. Palaearctic species not shown in Korea up to the present; relatively frequent but not numerous, occurs in various biotopes.

***Onthophagus (Onthophagus) bivertex* (HEYDEN, 1887)**

(Figs. 28, 29)

Distribution. USSR — Primorskij Kraj (Vladivostok), Ussuri; North China Korea, Japan (BALTHASAR, 1963).

Material examined. 179 ♂♂ and ♀♀. Prov. Phjŏngjang-si, Jongak-san (Mts, 40), 24 V 1964, M. MROCZKOWSKI and A. RIEDEL (ZIW), 29 V 1974, Z. STEBNICKA (ISEZ), Ponghva-ri (88), 24 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW); Prov. Phjŏngan-namdo, distr. Kangsŏ (61), vicinity of the lake Thesŏng, 26 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), 13 VI 1974, Z. STEBNICKA (ISEZ), distr. Sunan (104), Sŏkam-Čŏsudži (105), 30 VI 1974, Z. STEBNICKA (ISEZ), Junha-ri, 13 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW); Prov. Hamgjŏng-namdo, distr. Hongvŏn (30), Jŏnpong-ri, 8 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Hamgjŏng-pukto, distr. Kjŏngsŏng (44), Onpho-ri (77), 3 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Janggang-do, distr. Hjesan (26), Dŏngha-ri, 2 IX 1971, J. PAWŁOWSKI (ISEZ); Prov. Kangvŏn-do, Kymgang-san (Mts, 53), 16 VI 1974, Z. STEBNICKA (ISEZ); Prov. Kesŏng-si, vicinity of the water-fall Pakjŏn on the Čhŏnma-san (Mt., 13), 7 VI 1974, Z. STEBNICKA (ISEZ); Prov. Hvanghe-namdo, Sujang-san (Mt., 121), near Hedžu (18), 2 VI 1974, Z. STEBNICKA (ISEZ); Prov. Čhŏngdžin-si, valley of river Susŏng-čhŏn (110), 22 V 1974, Z. STEBNICKA (ISEZ).

Remarks. The species occurs in the pastures and other open areas in cow dung and at human excrements and is often collected.

***Onthophagus (Onthophagus) rugulosus* HAROLD, 1886**

Syn.: *Onthophagus sonani* MIWA, 1930.

(Figs. 33, 34)

Distribution. India (Assam), China (Yunnan, Sichuan, Fukien), Korea, Indochina — Vietnam (Tonkin), Taiwan (Nanshanchi, Funchiifo) (BALTHASAR, 1963; NOMURA, 1973).

Material examined. 35 ♂♂ and ♀♀. Prov. Hvanghe-namdo, vicinity of the water-fall Sujang-san (121) near Hedžu (18), 2—3 VI 1974, at human excrements, Z. STEBNICKA (ISEZ), valley of river Hakhjŏn-čhŏn, 3 VI 1974, in cow dung, Z. STEBNICKA (ISEZ); Prov. Kesŏng-si, vicinity of the water-fall Pakjŏn about 20 km SE of Kesŏng (56), 7 VI 1974, in sheep dung, Z. STEBNICKA (ISEZ); Prov. Phjŏngan-namdo, distr. Sunan (104), Sŏkam-Čŏsudži (105), 30 V 1974,

in cow dung on the banks of the lake, Z. STEBNICKA (ISEZ); Prov. Kangvön-do, Kymgang-san (Mts, 53) near Kosöng (122), 16 VI 1974, at human excrements, Z. STEBNICKA (ISEZ).

Remarks. Oriental species, relatively frequent in the southern Provinces of D.P.R.K.; collected mainly under human excrements, deep up to 30 cm below soil surface.

***Onthophagus (Onthophagus) olsoufieffi* BOUCOMONT, 1924**

Syn.: *Onthophagus uedanus* MATSUMURA, 1937.

(Figs. 30, 31)

Distribution. East Siberia (Ussuri), Manchuria, North China, Korea, Japan — Honshu (BALTHASAR, 1963).

Material examined. 11 ♂♂ and ♀♀. Prov. Hamgjöng-pukto, distr. Kjöngsöng (44), Onpho-ri (77), 21 VIII 1959, B. PISARSKI and J. PRÓSZYŃSKI (ZIW); Prov. Phjöngjang-si, Phjöngjang (86), 10 VI 1974, at human excrements, Z. STEBNICKA (ISEZ); Prov. Hvanghe-namdo, vicinity of the water-fall Sujang-san (121), 2 VI 1974, at human excrements, Z. STEBNICKA (ISEZ).

Remarks. Species found most frequently in human excrements or in the vicinity of them in the upper layer of soil.

***Onthophagus (Onthophagus) punctator* REITTER, 1892**

Syn.: *Onthophagus sunanicus* STEBNICKA, 1973, syn. n.

(Fig. 32)

Distribution. North China, Manchuria, Korea — Prov. Phjöngan-namdo, distr. Sunan (104), Sökam-Čösudži (105) (BALTHASAR, 1963; STEBNICKA, 1973).

Material examined. 35 ♂♂ and ♀♀. Prov. Kesöng-si, vicinity of the water-fall Pakjön on the Čhönma-san (Mt., 13), 6 VI 1974, Z. STEBNICKA (ISEZ); Prov. Čhöngdžin-si, valley of river Susöng-čhön (110), 22 V 1974, Z. STEBNICKA (ISEZ); Prov. Hvanghe-pukto, Nungni (132), 7 VIII 1950, BORCHSENIUS (ZIL); Prov. Hvanghe-namdo, vicinity of the water-fall Sujang-san (121), 2 VI 1974, Z. STEBNICKA (ISEZ).

Remarks. The taxonomic error made by the author (STEBNICKA, 1973) was due to the fact that in the descriptions of this species (REITTER, 1892: 179; BALTHASAR, 1963: 233, 489) the most characteristic feature, i. e. the shape of male anterior tibia had been omitted. BALTHASAR (1963) emphasises similar features in the descriptions of the following species: *O. atricapillus* D'ORB. and *O. nigellus* (ILLIG.). *O. punctator* REITT. is frequent and numerous in D.P.R.K.; most frequently found in the vicinity of waters in sandy soil under cow and sheep dung and at human excrements.

*Aphodiinae**Aphodiini**Aphodius (Colobopterus) apicalis* HAROLD, 1861

Syn.: *Aphodius haroldianus* BALTHASAR, 1932.

(Figs. 35—37)

Distribution. USSR — Transbaikalian Region, Ussuri, Primorskiy Kraj, Sakhalin, Kurile Islands (Alechino, Golovinno), North China, Manchuria, Korea, Japan — Hokkaido, Honshu (Murasugi, Matsunoyama), Sado (Mt. Kinpoku, Mt. Donden), Shikoku, Kyushu (NAKANE et BABA, 1960; BALTHASAR, 1963; PETROVITZ, 1972; KRYVOLUCKAJA, 1973; STEBNICKA, 1978).

Material examined. 62 ♂♂ and ♀♀. Prov. Kesöng-si, Čhönma-san (Mt., 13), 8 VI 1974, in cow dung, Z. STEBNICKA (ISEZ); Prov. Phjöngan-namdo, distr. Sunan (104), Sökam-Čösudži (105), 30 V 1974, in cow dung on the banks of the lake, Z. STEBNICKA (ISEZ), distr. Kangsö (61), Thesöng, 26 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), 13 VI 1974, Z. STEBNICKA (ISEZ); Prov. Phjöngjang-si, Tesöng-san (Mts, 124), 17 V 1974, Z. STEBNICKA (ISEZ); Prov. Janggang-do, distr. Hjesan (26), vicinity of the village Dönggha-ri, 2 IX 1971, J. PAWŁOWSKI (ISEZ); Prov. Čhöngdžin-si, Purjöng (87), valley of river, 23 V 1974, Z. STEBNICKA (ISEZ); Prov. Hvanghe-namdo, vicinity of the water-fall Sujang-san (121) near Hedžu (18), 2 VI 1974, Z. STEBNICKA (ISEZ), valley of river Hakhjön-čhön, 3 VI 1974, in cow dung (imagines and larvae of second and third stage) Z. STEBNICKA (ISEZ).

Description of larva: STEBNICKA, 1978.

Aphodius (Colobopterus) indagator MANNERHEIM, 1849

(Figs. 38—40)

Distribution. USSR — Transbaikalian Region (Irkutsk), East Siberia (BALTHASAR, 1964; PETROVITZ, 1972).


Material examined. 5 ♂♂ and ♀♀. Prov. Phjöngjang-si, Phjöngjang (86), 16 V 1974, Z. STEBNICKA (ISEZ); Prov. Hamgjöng-namdo, distr. Hjangsan (27), valley of river Hakhjön-čhön in the Mjohjang-san (Mts), 18 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Kangvön-do, Kymgang-san (Mts, 53) near Kosöng (122), 18 VI 1974, at human excrements, Z. STEBNICKA (ISEZ).

Remarks. This species is very similar to the former one: its distribution is little known. New to the Korean fauna, occurs sporadically and not frequently.

Aphodius (Colobopterus) subterraneus (LINNAEUS, 1758)

(Figs. 41, 42)

Distribution. Europe, North Africa (Algeria), Transcaucasus, Central Asia, Afghanistan, Mongolia, Siberia, Manchuria., North America (introduced) (BALTHASAR, 1964, ENDRÖDI, 1967; MEDVEDEV, 1974; STEBNICKA, 1976).



Figs. 35—37. *Aphodius (Colobopterus) apicalis* HAR. 35, 36 — male genitalia, lateral and dorsal view; 37 — female genitalia, stylus.

Figs. 38—40. *A. (C.) indagator* MANNRH.; 38 — female genitalia, stylus; 39, 40 — male genitalia, lateral and dorsal view. Figs. 41—44. Male genitalia, lateral and dorsal view; 41, 42 — *A. (C.) subterraneus* (L.); 43, 44 — *A. (Teuchestes) brachysomus* SOLS.

Material examined. 8 ♂♂ and ♀♀. Prov. Phjǽngjang-si, Jongak-san (Mt., 40), 28 V 1974, in sheep dung, Z. STEBNICKA (ISEZ); Prov. Čhǽngdžin-si, valley of river Susǽng-čhǽn (110), 25 V 1974, in cow dung, Z. STEBNICKA (ISEZ); Prov. Hamgǽng-pukto, distr. Kjǽngsǽng (44), valley Onpho-ri (77), 28 VIII 1959, B. PISARSKI and J. PRÓSZYŃSKI (ZIW).

Remarks. Palaearctic species, not shown in Korea up to now. Occurs mainly in lowlands and open pastures in cattle and sheep dung, at human excrements and in vegetable-detritus. Enemy of cultivation of fungi. A typical black form occurs in D.P.R.K. and also a rare variety ab. *fuscipennis* MULS. (elytra red); no numerous population found in Korea.

***Aphodius (Teuchestes) brachysomus* SOLSKY, 1874**

(Figs. 43, 44)

Distribution. USSR (Ussuri), China, Korea, Japan — Honshu, Kyushu, Takarajima (BALTHASAR, 1964; NOMURA, 1966).

Material examined. 21 ♂♂ and ♀♀. USSR — Primorskij Kraj, Chanka Lake, A. ZOLOTAREV (ZIL); Korea — Prov. Phjǽngan-namdo, distr. Kangsǽ, (61), 26 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Čhǽngdžin-si, distr. Purjǽng (87), 21 V 1974, in cow dung, Z. STEBNICKA (ISEZ), valley of river Susǽng-čhǽn (110), 22 V 1974, in cow dung, Z. STEBNICKA (ISEZ); Prov. Kesǽng-si, Kesǽng (56), 5 VI 1974, in sheep dung, Z. STEBNICKA (ISEZ).

Remarks. This species occurs sporadically and not frequently in D.P.R.K.; typical form only found.

***Aphodius (Teuchestes) haemorrhoidalis* (LINNAEUS, 1758)**

(Figs. 49, 50)

Distribution. Europe, Caucasus, Central Asia, Tibet, Mongolia, Siberia, Japan, North America (introduced) (NAKANE, 1960; BALTHASAR, 1964; ENDRÖDI, 1965; 1967; 1968; 1971; STEBNICKA, 1976).

Material examined. 110 ♂♂ and ♀♀. Prov. Janggang-do, distr. Hjesan (26), Dǽngha-ri, 2 IX 1971, J. PAWŁOWSKI (ISEZ); Prov. Čhǽngdžin-si, distr. Purjǽng (87), 21 V 1974, in cow dung, Z. STEBNICKA (ISEZ), valley of river Susǽng-čhǽn (110), 22 V 1974, in cow dung, Z. STEBNICKA (ISEZ); Prov. Hamgǽng-pukto, Kvanmo-bong (Mt., 60), 25 V 1974, at human excrements, Z. STEBNICKA (ISEZ), distr. Kjǽngsǽng (44), Džuyr (17), 25 V 1974, Z. STEBNICKA (ISEZ), Onpho-ri (77), 21 VIII 1959, B. PISARSKI and J. PRÓSZYŃSKI (ZIW); Prov. Kangvǽn-do, Kymgang-san (Mts, 53) near Kosǽng (122), 16 VI 1974, in cow dung, Z. STEBNICKA (ISEZ).

Remarks. Ubiquistic, very frequent and numerous species in the whole area of distribution. A typical form occurs in D.P.R.K., and also a very numerous variety ab. *sanguinolentus* Herbst. New to the Korean fauna.

Aphodius (Teuchestes) donghariensis STEBNICKA, 1973

(Fig. 48)

Distribution. D.P.R.K., Prov. Janggang-do, distr. Hjesan (26), Dõngha-ri (STEBNICKA, 1973).

Aphodius (Pharaphodius) rugosostriatus WATERHOUSE, 1875

(Figs. 45—47)

Distribution. North- and Central China, Korea, Japan — Hokkaido, Honshu, Sado (Hatano), Shikoku, Kyushu (NAKANE et BABA, 1960; NAKANE, 1960; BALTHASAR, 1964).

Material examined. 11 ♂♂ and ♀♀. Japan — Hokkaido, Kotyohama, 23 VII 1962, Y. SHIBATA (ISEZ); Korea — Prov. Hvanghe-pukto, Sarivõn (101), 20 VII 1950, BORCHSENIUS (ZIL), 10 VII 1956, M. MAGYAR (HMNH); Prov. Phjõngan-namdo, distr. Kangsõ (61), Thesõng, 24 VIII 1971, J. RAZOWSKI (ISEZ).

Remarks. Species very similar to *A. (Coptochiroides) subcostatus* KOLBE. In Korea occurs sporadically only in summer; probably gives one generation per year.

Aphodius (Coptochiroides) subcostatus KOLBE, 1886

(Figs. 51, 52)

Distribution. China, Korea, Japan, Taiwan, North Vietnam (BALTHASAR, 1964).

Material examined. 4 ♂♂ and ♀♀. Prov. Hvanghe-pukto, Sarivõn (101), 2—20 VII 1950, BORCHSENIUS (ZIL).

Remarks. Oriental species, similarly as *A. (Pharaphodius) rugosostriatus* WATERH. is rare and not frequent in Korea.


Aphodius (Acrossus) rufipes (LINNAEUS, 1758)

Syn.: *Aphodius matsuzawai* YAWATA, 1942.

(Figs. 53, 54)

Distribution. Europe, Transcaucasus, Iran, Siberia, Kurile Islands, Japan — Honshu (Utsukushigahara, Mt. Yatsugatake, Mt. Tateshima, Mt. Asama, Shinshu-toge), North- and South America (introduced), South Africa (introduced) (BALTHASAR, 1964; NAKANE et MASUMOTO, 1967; MEDVEDEV et ERMOLENKO, 1969; KRYVOLUCKAJA, 1973; STEBNICKA, 1976).

Material examined. 3 ♂♂. Prov. Hamgjõng-pukto, distr. Kjõngsõng (44), Onpho-ri (77), 11 IX 1966, H. SZEŁĘGIEWICZ and C. DZIADOSZ (ZIW); Prov.



Figs. 45—47. *Aphodius (Pharaphodius) rugosostriatus* WATERH. 45, 46 — male genitalia, lateral and dorsal view; 47 — female genitalia, stylus.

Fig. 48. *A. (Teuchestes) donghariensis* STEB. — female head. Figs. 49—54. Male genitalia, lateral and dorsal view; 49, 50 — *A. (T.) haemorrhoidalis* (L.); 51, 52 — *A. (Coptochiroides) subcostatus* KOLBE; 53, 54 — *A. (Acrossus) rufipes* (L.). Figs. 55—57. *A. (A.) binaevulus* HEYD. 55 — female genitalia, stylus; 56, 57 — male genitalia, lateral and dorsal view

Phjôngjang-si, Jongak-san (Mt., 40), 29 V 1974, in cow dung, Z. STEBNICKA (ISEZ).

Remarks. The Korean specimens do not differ significantly from the va-

riable individuals of this species known as very common and numerous one in Europe; in D.P.R.K. it occurs seldom, exclusively in the mountains. New to the Korean fauna.

Aphodius (Acrossus) binaevulus HEYDEN, 1887

(Figs. 55—57)

Distribution. USSR — Primorskij Kraj (Vladivostok), Ussuri (BALTHASAR, 1964).

Material examined. 30 ♂♂ and ♀♀. Prov. Phjŏngjang-si, Jongak-san (Mt., 40), 29 V 1974, in cow dung, Z. STEBNICKA (ISEZ); Prov. Hamgjŏng-pukto, Kvanmo-bong (Mt., 60), 23 V 1974, at human excrements, Z. STEBNICKA (ISEZ); Prov. Janggang-do, distr. Samdžijŏn (103), Namphote-san (76), 8 IX 1971, J. PAWŁOWSKI (ISEZ); Prov. Čhŏngdžin-si, Purjŏng (87), 24 V 1974, in sheep dung, Z. STEBNICKA (ISEZ).

Remarks. New species to the Korean fauna, occurs in the mountains. Several coloured variants found; most frequently is represented the variety *ab. diaphanomaculatus* HEYD. (head and pronotum black, elytra red), less numerous is the variety *ab. niger* TES. (whole body black), a typical form is seldom (elytra black with yellowish or reddish spots near the apex). This species is very similar to the Euro-Siberian one *A. (A.) depressus* (KUG.) whose area extends as far as to Bajkal. It seems that *A. binaevulus* HEYD. replaces *A. depressus* in the same habitat to the south-east of the latter's range.

Aphodius (Acrossus) superatratus NOMURA et NAKANE, 1951

(Figs. 58, 59)

Distribution. Japan — Honshu (Buhei-toge, Kyoto, Kibune, Mt. Daibosatsu, Nishi-Tanzawa) (BALTHASAR, 1964; NAKANE et MASUMOTO, 1967).

Material examined. 13 ♂♂ and ♀♀. Prov. Hamgjŏng-pukto, Kvanmo-bong (Mt., 60), 23 V 1974, at human excrements, Z. STEBNICKA (ISEZ).

Description of the Korean specimens. Length 6—7.5 mm. Elongate-oval, rather strongly convex, shining, black. Clypeal margin finely reflexed, broadly rounded each side of very shallow median emargination, sides arcuate to sharply rounded, right-angled genae; surface of head with rather evenly spaced, moderately coarse punctures, generally separated by their diameters over middle disc, finer and closer anteriorly and a little smaller basally. Pronotum moderately convex, sides strongly margined, base without marginal line; surface very minutely alutaceus throughout under high magnification, mixed moderately fine and coarse, quite evenly distributed punctures throughout except for a narrow, impunctate, longitudinal midline; the punctures are larger and closer laterally. Scutellum triangular with a few coarse punctures at base. Elytra convex, oval, humeri distinctly dentate, striae moderately deep, very

finely and indistinctly punctate; intervals slightly convex with moderately fine, close, evenly distributed punctures which practically contiguous the same size as finer punctures of the pronotum; lateral and apical part of elytra shortly piliferous. Ventral surface and femora alutaceous, closely punctate, metasternum flat, midline shallow, metasternal surface distinctly punctate. Apical spur of fore tibia straight and acute; apex of middle and hind tibiae fringed with moderately long, unequal spines; first posterior tarsal segment noticeably shorter than the upper spur, equal to the remaining segments combined.

Male. Pronotum more convex than in female. Aedeagus as in figures 58, 59.

Female. Pronotum less convex than in male.

Remarks. Little known species; as the majority of representatives of the subgenus *Acrossus* MULS. it occurs exclusively in the mountains.

***Aphodius (Paulianellus) maderi* BALTHASAR, 1938**

Syn.: *Aphodius asahinai* NAKANE, 1951.

(Figs. 60, 61)

Distribution. China (Sichuan), Japan — Honshu (Nara), Kyushu (BALTHASAR, 1964; NAKANE et MASUMOTO, 1967).

Material examined. 3 ♂♂. Prov. Kangvön-do, Kymgang-san (Mts, 53), vicinity of Kosöng (122), 18 VI 1974, Z. STEBNICKA (ISEZ).

Remarks. Little known species distinguished by the features given in the diagnosis of the subgenus *Paulianellus* BALTH. It is closely related to the representatives of subgenus *Acrossus* MULS. New to the Korean fauna.

***Aphodius (Pleuraphodius) lewisi* WATERHOUSE, 1875**

(Fig. 64)

Distribution. Ceylon, Java, Lombok, Philippines, India (Assam), China, Taiwan (Liyutan, Kenting Park), Japan — Honshu, Kyushu, Hachijo-Kojima; Korea (BALTHASAR, 1964; NOMURA, 1969; 1973).

Material examined. 42 ♂♂ and ♀♀. Prov. Kesöng-si, Pakjön 20 km SE of Kesöng (56), 8 VI 1970, S. MAHUNKA and H. STEINMANN (HMNH), 8 VI 1974, at human excrements, Z. STEBNICKA (ISEZ); Prov. Phjöngan-namdo, distr. Sunčhön (94), Džamo-san, 27 VIII 1971, at light of UV lamps, J. PAWŁOWSKI (ISEZ); Prov. Kangvön-do, Kymgang-san (Mts, 53), 18 VI 1974, in cow dung, Z. STEBNICKA (ISEZ); Prov. Phjöngjang-si, Jongak-san (Mt., 40), 20 VIII 1971, at light of UV lamps, J. PAWŁOWSKI (ISEZ).

Description of the Korean specimens. Length 2—3 mm. Elongate-oblong, moderately convex, shining, dark reddish-brown or brown, posterior part of head and disc of pronotum sometimes darker. Head trapezoid, moderately convex without tubercles, frontal suture feebly marked; clypeal margin finely reflexed, obtusely rounded each side of moderate median emargination; sur-

face punctures fine, evenly distributed, slightly alutaceous within. Pronotum convex, sides slightly arcuate, anterior and posterior angles obtusely rounded, sides finely margined, base with a row of dense, fine punctures, without marginal line; surface alutaceous, densely, moderately coarsely punctate throughout. Scutellum triangular, narrow, impunctate. Elytra oval, humeri not dentate; striae moderately deep, shining, closely and finely punctate, the punctures not crenating inner margins of the intervals, the 8th stria long, approaching the base of elytra; intervals moderately wide, not connected apically, each interval laterally flattened, mat with shining, impunctate carina at middle. Metasternum shining, finely and densely punctate, midline distinctly but not deeply impressed; abdomen shining, finely punctate. Legs slender; apical spur of fore tibia straight, acute; middle and hind tibiae thin, apical fringe with thin setae unequal in length; first posterior tarsal segment long, about half time longer than the upper spur, slightly longer than following three segments combined.

Male. The last abdominal segment slightly shorter than in female. Aedeagus as in figure 64.

Female. The last abdominal segment slightly longer than in male.

Remarks. The subgenus *Pleuraphodius* A. SCHM. contains about 22 species, that inhabit mainly the Ethiopian Region, Oriental Realm and East Palaearctic. *P. lewisi* WATERH. is considered as a rare species (BALTHASAR, 1964); it is relatively frequent but not numerous in the southern Provinces of D.P.R.K.

Aphodius (Trichaphodius) proclivis BALTHASAR, 1933

(Figs. 62, 63)


Aphodius (Gilletianus) proclivis BALTHASAR, 1933:139 (description of male); BALTHASAR, 1964: 177—178 (redescription).

Distribution: South China (Yunnan-Sen).

Material examined. 95 ♂♂ and ♀♀. Prov. Hamgjông-namdo, distr. Hjangsan (27), Hjangam-ri, 18 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Kesông-si, vicinity of the water-fall Pakjôn on the Čhônma-san (Mt., 13), 6—8 VI 1974, numerous specimens found under cow dung and at human excrements, Z. STEBNICKA (ISEZ); Prov. Hvanghe-namdo, vicinity of the water-fall Sujang-san (121) near Hedžu (18), 2 VI 1974, Z. STEBNICKA (ISEZ), valley of river Hakhjôn-čhôn below Sujang-san (121) water-fall, 3 VI 1974, in cow dung, Z. STEBNICKA (ISEZ); Prov. Kangvôn-do, Kymgang-san (Mts, 53), 18 VI 1974, in cow dung, Z. STEBNICKA (ISEZ).

Description of the Korean specimens. Length 4.3—5.5 mm. Oblong-elongate, moderately convex, shining, dark brown with sides of pronotum, legs and usually base and apical part of elytra partially yellowish. Antennae light yellow, antennal clubs brownish, terminal segment of maxillary palpus cylindrical. Head

broad, very weakly convex, strongly reflexed clypeal margin broadly rounded each side of very shallow median emargination, sides arcuate to sharply rounded right-angled genae; surface very minutely alutaceous with rather evenly spaced, fine punctures. Pronotum rather weakly convex, sides moderately arcuate, anterior angles widely rounded, posterior angles obtuse, sides and base



Figs. 58—68. Male genitalia, lateral and dorsal view; 58, 59 — *Aphodius (Acrossus) superatrus* NOM. et NAK.; 60, 61 — *A. (Paulianellus) maderi* BALTH.; 62, 63 — *A. (Trichaphodius) proclivis* BALTH.; 64 — *A. (Pleuraphodius) lewisi* WATERH.; 65, 66 — *A. (Aganocrossus) urostigma* HAR.; 67, 68 — *A. (Trichonotulus) mongolicus* MANNRH. Fig. 69. *A. (Aphodius) elegans* ALLIB. — left elytron

finely margined; surface very minutely alutaceous throughout under high magnification, mixed very fine and fine, unevenly distributed punctures a trifle smaller on the disc, larger and closer laterally. Scutellum triangular, narrow, finely punctate. Elytra convex, subparallel with sides and apical part shortly,

distinctly piliferous; striae fine, not deep, very fine strial punctures weakly crenating inner margins of the moderately convex, minutely alutaceous intervals; intervals with very fine punctures, closer and more distinct at apex of elytra. Metasternum minutely alutaceous, impunctate, midline weakly impressed, abdominal sterna alutaceous, finely punctate. Anterior tibia with three lateral teeth, crenate toward the base, the terminal spur straight; middle and hind tibiae long, slender, apical fringe with rather long setae, distinctly unequal in length; first posterior tarsal segment very long, about one-third time longer than the upper spur and clearly longer than following three segments combined.

Male. Apical spur of fore tibia wider than in female, truncate at apex; metasternum slightly concave, pronotal punctures less coarse. Aedeagus as in figures 62, 63.

Female. Apical spur of fore tibia narrower than in male, sharply pointed; metasternum not concave, pronotal punctures more coarse.

Remarks. I have examined the holotype of *A. proclivis* in the BALTHASAR's collection in Prague. This species has been described by BALTHASAR (1933) in a new subgenus *Gilletianus* BALTH. The establishment of this subgenus was not justified, since the complex of the features of this species indicates its affiliation to the subgenus *Trichaphodius* A. SCHM. New to the Korean fauna, frequent and numerous in various biotopes.

Aphodius (Aganocrossus) urostigma HAROLD, 1862

(Figs. 65, 66)

Distribution. Sunda Islands (Java, Borneo, Sumatra, Celebes), Indochina (Malaya, Thailand, Laos, Vietnam, Burma), India (Assam), Nepal, China (Hainan, Yunnan), South Afghanistan, Taiwan (Liyutan, Nanshanchi, Liukuei Lutaol Isl., Lanyu Isl.), Korea, Japan — Hokkaido, Honshu, Shikoku, Kyushu, Yakushima, Kuchinoerabu, Nakanoshima, Takarajima, Amami-Ōshima, Iki, Tsushima, Ishigaki, Iriomote, Ohshima, Toshima, Miyake, Hachijo, Hachijo-Kojima, Aogashima, Tokara; Africa (introduced) (BALTHASAR, 1964; 1965; NOMURA, 1966; 1969; 1973).

Material examined. 95 ♂♂ and ♀♀. Japan — Honshu, Kanagawa Pref., Sagami-hara, 13 VII 1967, Y. SHIBATA (ISEZ), Yamanashi Pref., near Gozaishi, 27 VII 1965, Y. SHIBATA (ISEZ), Tokyo, Machida, Tsurukawa, 6 VIII 1968, Y. SHIBATA (ISEZ); Korea — Prov. Hvanghe-namdo, Sinchŏn (108), 18 IX 1971, J. PAWŁOWSKI (ISEZ), vicinity of the water-fall Sujang-san (Mt., 121) near Hedžu (18), 1—3 VI 1974, Z. STEBNICKA (ISEZ); Prov. Kangvŏn-do, Kymgang-san (Mts, 53), 16—18 VI 1974, in cow dung and at human excrements, Z. STEBNICKA (ISEZ); Prov. Phjŏngan-namdo, distr. Kangsŏ (61), Thesŏng, 13 VI 1974, in cow dung, Z. STEBNICKA, 25 VIII 1971, J. RAZOWSKI (ISEZ), distr. Sunan (104), Sŏkam-Čŏsudži (105), 30 V 1974, in cow dung, Z. STEB-

NICKA (ISEZ), distr. Sunčhön (94), Džamo-san, 28 VIII 1971, at light of UV lamps, J. RAZOWSKI (ISEZ); Prov. Phjǽngjang-si, Phjǽngjang (86), 15 IX 1971, at human excrements, J. PAWŁOWSKI (ISEZ), Ponghva-ri (88), 24 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW); Prov. Hamgjǽng-namdo, distr. Hjangsan (27), Hjangam-ri, Mjohjang-san (Mts), 18 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. Species frequent and numerous in D.P.R.K., occurs in various biotopes.

***Aphodius (Trichonotulus) dzamosanicus* STEBNICKA, 1973**

(Fig. 70)

Distribution. D.P.R.K. Prov. Phjǽngan-namdo, distr. Sunčhön (94), Džamo-san (STEBNICKA, 1973).

***Aphodius (Trichonotulus) mongolicus* MANNERHEIM, 1852**

(Figs. 67, 68)

Distribution. East Siberia, Mongolia (BALTHASAR, 1965; ENDRÖDI, 1971).

Material examined. 5 ♂♂ and ♀♀. Prov. Čhǽngdžin-si, valley of river Susǽng-čhön (110), 21 V 1974, in sheep dung, Z. STEBNICKA (ISEZ).

Remarks. Species very similar to the Euro-Caucasian one *A. (T.) scrofa* (F.); slightly differs from it by shape and punctuation of the head. According to BALTHASAR (1964) it is probably a subspecies of *A. (T.) scrofa*, however, its distribution is little known as yet.

***Aphodius (Aphodius) elegans* ALLIBERT, 1847**

(Fig. 69)

Distribution. China, Japan — Hokkaido, Honshu, Sado (Kitauzima, Kawaharada, Mt. Kinpoku, Ookura-goe), Shikoku, Kyushu, Takarajima, Amami-Ōshima, Miyake, Okinawa; Taiwan, Korea, Indochina (NAKANE et BABA, 1960; BALTHASAR, 1964; NOMURA, 1966; 1969).

Material examined. 2 ♂♂. Prov. Hamhyng-si, Hyngnam (28), 7 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Phjǽngan-namdo, distr. Sunan (104), Junha-ri, 3 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW).

***Aphodius (Orodalus) pusillus* (HERBST, 1789)**

(Figs. 71, 72)

Distribution. Europe, Transcaucasus, Siberia, Mongolia, Japan — Hokkaido, Honshu (Niigata, Shibata), Shikoku, Kyushu (NAKANE, 1960; NAKANE et BABA, 1960; BALTHASAR, 1964; ENDRÖDI, 1971 et all.).

Material examined. 82 ♂♂ and ♀♀. Japan — Honshu, Kanagawa Pref., Atsugi, 7 VI 1960, Y. SHIBATA (ISEZ), Mt. Oogusu, 3 VI 1961, Y. SHIBATA (ISEZ), Nagano Pref. Utsukushi, 1 V 1960, Y. SHIBATA (ISEZ); Korea — Prov. Chŏngdžin-si, valley of river Susŏng-čhŏn (110), 22 V 1974, numerous specimens in cow dung, Z. STEBNICKA (ISEZ), Purjŏng (87), valley of river, 22 V 1974, in sheep dung, Z. STEBNICKA (ISEZ); Prov. Phjŏngjang-si, Jongak-san (Mt., 40), 29 V 1974, Z. STEBNICKA (ISEZ); Prov. Kangvŏn-do, Kymgang-san (Mts, 53), 16 VI 1974, in cow dung, Z. STEBNICKA (ISEZ).

Remarks. In the whole distributional area this species indicates great individual variability; it mainly concerns punctation of head and pronotum, sculpture of elytra, color and body size. The distinction of Far-Eastern subspecies based on the small differences in external morphology by some authors is not sufficiently justified. Species new to the Korean fauna, occurs in some color variants. Locally numerous, collected mainly in cow and sheep dung.

Aphodius (Orodalus) naraensis NAKANE, 1956

(Figs. 73—75)

Distribution. Japan — Honshu (Nara) (NAKANE, 1960; BALTHASAR, 1964).

Material examined. 9 ♂♂ and ♀♀. Prov. Chŏngdžin-si, valley of river Susŏng-čhŏn (110), 22 V 1974, in cow dung, Z. STEBNICKA (ISEZ); Prov. Kesŏng-si, vicinity of the water-fall Pakjŏn on the Čhŏnma-san (Mt., 13), 7 VI 1974, in sheep dung, Z. STEBNICKA (ISEZ).

Description of the Korean specimens. Length 3—3.5 mm. Elongate oblong, moderately convex, black; margin of clypeus, shoulders and apex of elytra reddish, sometimes disc of elytra reddish brown. Head slightly convex, finely reflexed clypeal margin obtusely rounded each side of moderately deep anterior emargination, sides weakly arcuate to small, slightly flattened genae; frontal suture distinctly marked, surface of head everywhere closely, uniformly, moderately punctate, the punctures generally separated by less than their diameters. Pronotum more or less convex, slightly alutaceous, posterior angles obtusely rounded, sides and base margined; surface everywhere very densely punctate, the punctures moderate in size, very slightly smaller and closer toward the sides. Scutellum elongate, slightly convex at apex, finely punctate. Elytra subparallel, striae moderately deep, strial punctures as wide as striae, distinctly crenating edges of the intervals; intervals more or less convex, alutaceous with close, fine punctures. Ventral surface alutaceous, finely punctate, metasternum with moderately deep midline. Apex of middle and hind tibiae fringed with short, unequal spines; first segment of posterior tarsus longer than the upper spur and nearly equal to the following two segments combined.

Male. Apical spur of fore tibia bent inward at apex; aedeagus as in figures 74, 75.

Female. Apical spur of fore tibia straight; stylus as in figure 73.

Remarks. The original description of the male holotype of *A. naraensis* is incomplete. The holotype was not examined by the author of the present study. Series of specimens from Korea is characterized by great individual variability; this concerns punctuation and color of elytra and shape of pronotum. Body shape of this species is very similar to this of *A. pusillus* (HERBST); both species occur in the same biotopes. New to the Korean fauna.




Fig. 70. *Aphodius* (*Trichonotulus*) *dzamosanicus* STEB. — female head.
 Figs. 71, 72. *A. (Orodalus) pusillus* (HBST) — male genitalia, lateral and dorsal view. Figs. 73—75. *A. (O.) naraensis* NAK. 73 — female genitalia, stylus; 74, 75 — male genitalia, lateral and dorsal view.
 Figs. 76—82. Male genitalia, lateral and dorsal view. 76, 77 — *A. (Agrilinus) uniformis* WATERH.; 78, 79 — *A. (A.) brevisculus* (MOTSCH.); 80 — *A. (Phaeaphodius) rectus* MOTSCH.; 81, 82 — *A. (Agrilinus) putridus* (HBST)

Aphodius (Phaeaphodius) rectus MOTSCHULSKY, 1866

(Fig. 80)

Distribution. Central Asia (Iran), Mongolia, East Siberia (Bajkal Region), Sachalin, Kurile Islands (Southern Kurilsk, Alechino, basin of Kurilka river), Primorskij Kraj; North- and Central China, Korea, Japan — Hokkaido, Honshu, Shikoku, Kyushu, Sado, Ohshima, Miyake, Hachijo, Hachijo-Kojima; Taiwan (BALTHASAR, 1964; TESÁŘ, 1968; NOMURA, 1969; KRYVOLUCKAJA, 1973).

Material examined. 130 ♂♂ and ♀♀. USSR — Bajkal, Listvianka, 26—30 VI 1974, in cow dung, Z. STEBNICKA (ISEZ); Japan — Honshu, Yamanashi Pref., Nagasawa, 1 IV 1967, Y. SHIBATA (ISEZ), Kanagawa Pref., Mt. Oogusu, 3 IV 1961, Y. SHIBATA (ISEZ), Mt. Ooyama, 23 IX 1967, 25 XII 1966, Y. SHIBATA (ISEZ), Ibaraki Pref., Watarase, 4 V 1961, Y. SHIBATA (ISEZ), Tokyo Pref., Mt. Takao, 28 III 1960, Y. SHIBATA (ISEZ), Tamakyuryo, 18 III 1966, Y. SHIBATA (ISEZ), Setagaya, 10 II 1961, Y. SHIBATA (ISEZ); Korea — Prov. Janggang-do, distr. Hjesan (26), valley Dongha-ri, 2 IX 1971, J. PAWŁOWSKI (ISEZ); Prov. Čhôngdžin-si, valley of river Susông-čhôn (110), 21—25 V 1974, in cow and sheep dung, Z. STEBNICKA (ISEZ), Purjông (87), 21—25 V 1974, in cow dung, Z. STEBNICKA (ISEZ); Prov. Hamgjong-punkto, distr. Kjôngsông (44), valley Onpho-ri (77), 23 V 1974, in sheep dung, Z. STEBNICKA (ISEZ); Prov. Hvanghe-namdo, water-fall Sujang-san (121) near Hedžu (18), 2 VI 1974, Z. STEBNICKA (ISEZ), valley of river Hakhjôn-čhôn, 3 VI 1974, Z. STEBNICKA (ISEZ); Prov. Phjôngjang-si, Jongak-san (Mt., 40), 29 V 1974, in sheep dung, Z. STEBNICKA (ISEZ); Prov. Kangvôn-do, Kymgang-san (Mts, 53), 18 VI 1974, Z. STEBNICKA (ISEZ).

Remarks. This species is very characteristic representative of Manchurian fauna. In D.P.R.K. locally frequent, mainly in the northern Provinces. Typical black form occurs with the variety *ab. biformis* REITT.

Aphodius (Agrilinus) uniformis WATERHOUSE, 1875

Syn.: *Aphodius etorofuensis* KONÔ, 1937;

Aphodius maritimus NOMURA et NAKANE, 1951.


(Figs. 76, 77)

Distribution. Kurile Islands, Korea, Japan — Hokkaido, Honshu, Shikoku, Kyushu, Nakanoshima, Takarajima, Amami-Ōshima, Iriomote, Tsushima, Miyake, Hachijo, Aogashima (BALTHASAR, 1964; NOMURA, 1966; 1969; KRYVOLUCKAJA, 1973).

Material examined. 70 ♂♂ and ♀♀. Prov. Čhôngdžin-si, valley of river Susông-čhôn (110), 22 V 1974, in cow dung, Z. STEBNICKA (ISEZ), Purjông (87), 21 V 1974, Z. STEBNICKA (ISEZ); Prov. Hvanghe-namdo, vicinity of the water-fall Sujang-san (121) near Hedžu (18), 3 VI 1974, in sheep dung, Z. STEBNICKA (ISEZ); Prov. Kesông-si, vicinity of the water-fall Pakjôn, 20 km SE of Kesông

(56), 8 VI 1974, at human excrements, Z. STEBNICKA (ISEZ); Prov. Phjônggan-namdo, distr. Sunčhôn (94), Džamo-san, 27 VIII 1971, at light of UV lamps, J. RAZOWSKI (ISEZ), distr. Sunan (104), Sôkam-Čôsudži (105), 30 V 1974, in cow dung, Z. STEBNICKA (ISEZ), Junha-ri, 40 km W of Phjônggang (86), 13 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), distr. Kangsô (61), Thesông, 24 VIII 1971, at light of UV lamps, J. RAZOWSKI (ISEZ), 13 VI 1974, in cow dung, Z. STEBNICKA (ISEZ), 26 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. Frequent and relatively numerous species in D.P.R.K., occurs in various biotopes, in cattle dung and human excrements.



Figs. 83—86. Male genitalia, lateral and dorsal view; 83, 84 — *Aphodius (Bodilus) languidulus* A. SCHM.; 85, 86 — *A. (Calamosternus) uniopagiatus* WATERH. Figs. 87—89, *A. (Agrilinus) inexpectatus* BALTH. 87 — female genitalia, stylus; 88, 89 — male genitalia, lateral and dorsal view. Figs. 90—95. Male genitalia, lateral and dorsal view; 90, 91 — *A. (Bodilus) sordidus* (FABR.); 92, 93 — *A. (Calamosternus) sublimbatus* (MOTSCH.); 94, 95 — *A. (Nialus) sturmi* HAR.

Aphodius (Agrilinus) putridus (HERBST, 1789)

(Figs. 81, 82)

Distribution. North- and Central Europe, Caucasus, North Siberia, Mongolia, North America (introduced) (BALTHASAR, 1964; ENDRÖDI, 1967; 1971; STEBNICKA, 1976 et al.).

Material examined. 1 ♂ and 2 ♀♀. Prov. Janggang-do, distr. Samdžijon (103), vicinity of the village Posŏ-ri, 8 IX 1971, in cow dung, J. PAWŁOWSKI, (ISEZ).

Remarks. Euro-Siberian species, new to the Korean fauna. In D.P.R.K. variety *ab. transitus* found, which differs from the typical form by color of elytra (black with reddish spots on shoulders and at apex) and strongest punctation.

Aphodius (Agrilinus) brevisculus (MOTSCHULSKY, 1866)

Syn.: *Aphodius nigerrimus* WATERHOUSE, 1875.

(Figs. 78, 79)

Distribution. Kurile Islands (Kurilsk), Korea, Japan — Hokkaido, Honshu, Shikoku, Kyushu (NAKANE, 1961; BALTHASAR, 1964; KRYVOLUCKAJA, 1973).

Material examined. 72 ♂♂ and ♀♀. Prov. Čhŏngdžin-si, valley of river Su-sŏng-čhŏn (110), 22 V 1974, numerous specimens collected in cow dung, Z. STEBNICKA (ISEZ), Purjŏng (87), valley of river, 22 V 1974, Z. STEBNICKA (ISEZ).

Remarks. Probably local species, sometimes numerous, collected in D.P.R.K. only in one of northern Provinces.

Aphodius (Agrilinus) inexpectatus BALTHASAR, 1935

(Figs. 87—89)

Distribution. USSR — Primorskij Kraj (Vladivostok) (BALTHASAR, 1964).

Material examined. 5 ♂♂ and ♀♀. Prov. Kangvŏn-do, Kymgang-san (Mts, 53), 18 VI 1974, in sheep dung, Z. STEBNICKA (ISEZ); Prov. Hamgjŏng-pukto, distr. Kjŏngsŏng (44), valley Onpho-ri (77), 21 VIII 1959, B. PISARSKI and J. PRÓSZYŃSKI (ZIW).

Description of the Korean specimens. Length 4.5—5.5 mm. Oblong, subparallel, moderately convex, shining, dark red-brown, sometimes disc of pronotum and lateral part of elytra darker. Head moderately convex, edge finely reflexed, clypeus slightly angulate each side of moderate median emargination, sides nearly straight to small, obtusely rounded genae; surface of head finely punctate, frontal suture with three more or less distinct tubercles. Pronotum moderately convex, posterior angles obtusely rounded, sides straight or weakly emarginated over anterior three-fifths then weakly arcuate to posterior angles, side and base finely margined; surface densely punctate with mixed fine and

moderately coarse punctures, closer and larger laterally. Scutellum triangular, distinctly punctate. Elytra convex, sides weakly arcuate, humeri not dentate; striae deep, rather wide, strial punctures slightly crenating inner margins of the narrow, strongly convex, not connected at apex intervals; surface of intervals impunctate, shining. Ventral surface shining; metasternum finely punctate, midline shallow. Apical spur of fore tibia straight and acute; apex of middle and hind tibiae fringed with short, equal spines; first segment of posterior tarsus shorter than upper spur and slightly longer than the next two segments combined.

Male. Frontal tubercles distinctly marked, shorter spur of middle tibia slightly truncate at apex. Aedeagus as in figures 88, 89.

Female. Frontal tubercles less distinctly marked than in male, shorter spur of middle tibia sharply pointed. Stylus as in figure 87.

Remarks. Species new to the Korean fauna. The holotype (only one male specimen known) belongs to the private collection of V. BALTHASAR in Prague.

Aphodius (Bodilus) sordidus (FABRICIUS, 1775)

(Figs. 90, 91)

Distribution. North- and Central Europe, Transcaucasia, Siberia (Omsk), China (Yunnan), Korea, Japan — Hokkaido, Honshu, Shikoku, Kyushu, Oshima, Hachijo, Yakushima (BALTHASAR, 1964; TESAŘ, 1968; NOMURA, 1969; STEBNICKA, 1976; MEDVEDEV, 1976 et al.).

Material examined. 45 ♂♂ and ♀♀. Prov. Kesöng-si, Pakjön, 20 km SE of Kesöng (56), 8 VI 1970, S. MAHUNKA and H. STEINMANN (HMNH), vicinity of the water-fall Pakjön, 6—8 VI 1974, in sheep dung, Z. STEBNICKA (ISEZ); Prov. Hvanghe-namdo, vicinity of Sinchön (108), 16 IX 1971, J. PAWŁOWSKI (ISEZ), Sujang-san (Mt., 121) near Hedžu (18), 3 VI 1974, in cow dung, Z. STEBNICKA (ISEZ); Prov. Janggang-do, distr. Hjesan (26), valley Dongha-ri, 2 IX 1971, in cow dung, J. PAWŁOWSKI (ISEZ), Hjesan (26), 11 IX 1971, at light of UV lamps, J. RAZOWSKI (ISEZ); Prov. Hamgjöng-pukto, distr. Kjöngsöng (44), valley Onpho-ri (77), 21 VIII 1959, B. PISARSKI and J. PRÓSZYŃSKI (ZIW); Prov. Phjöngan-namdo, distr. Kangsö (61), Thesöng, 26 V 1965, M. MROCKOWSKI and A. RIEDEL (ZIW).

Remarks. Euro-Asiatic species; occurs in various biotopes, in particular in the regions rich in limestone. In the D.P.R.K. frequent but not very numerous.

Aphodius (Bodilus) languidulus A. SCHMIDT, 1922

(Figs. 83, 84)

Distribution. North- and East China, Korea, Japan — Honshu (Kurokawa), Shikoku, Kyushu (NAKANE et BABA, 1960; NAKANE, 1961; BALTHASAR, 1964).

Material examined. 9 ♂♂ and ♀♀. Prov. Phjöngan-namdo, distr. Kangsö

(61), Thesŏng, 24 VIII 1971, at light of UV lamps, J. RAZOWSKI (ISEZ); Prov. Janggang-do, distr. Hjesan (26), 1 IX 1971, at light of UV lamps, J. PAWŁOWSKI (ISEZ); Prov. Hvanghe-namdo, Sinčhŏn (108), 16 IX 1971, in cow dung, J. PAWŁOWSKI (ISEZ); Prov. Hvanghe-pukto, Sarivŏn (101), 19 VII 1950, BORCHSENIUS (ZIL).

Remarks. Not frequent and not numerous species. It is probably an element of the Manchurian fauna of narrow area of distribution. In Korea collected mainly at light.

Aphodius (Nialus) sturmi HAROLD, 1870

Syn.: *Aphodius inouei* NOMURA, 1942.

(Figs. 94, 95)

Distribution. South- and South-Eastern Europe, North Africa, Asia Minor, Central Asia, Mongolia, Manchuria, Korea, Japan (BALTHASAR, 1964; ENDRÖDI, 1965; MEDVEDEV, 1976).

Material examined. 120 ♂♂ and ♀♀. Prov. Hamgjŏng-pukto, distr. Kjŏngsŏng (44), valley Onpho-ri (77) 21 VIII 1959, B. PISARSKI and J. PRÓSZYŃSKI (ZIW); Prov. Phjŏngan-namdo, distr. Kangsŏ (61), Thesŏng, 24 VIII 1971, J. RAZOWSKI (ISEZ), distr. Sunčhŏn (94), Džamo-san, 27 VIII 1971, collected at light of UV lamps, J. PAWŁOWSKI (ISEZ), distr. Sunan (104), Sŏkam-Čŏsudži (105), 21 VIII 1971, at light, J. PAWŁOWSKI (ISEZ), 30 V 1974, in cow and sheep dung, Z. STEBNICKA (ISEZ); Prov. Phjŏngjang-si, Jongak-san (Mt., 40), 16—20 VIII 1971, at light of UV lamps, J. RAZOWSKI (ISEZ); Prov. Hvanghe-namdo, vicinity of Sinčhŏn (108), 16 IX 1971, in cow dung, J. PAWŁOWSKI (ISEZ), Sujang-san (Mt., 121) near Hedžu (18), 2 VI 1974, at human excrements, Z. STEBNICKA (ISEZ), valley of river Hakhjŏn-čhŏn, 3 VI 1974, in cow dung, Z. STEBNICKA (ISEZ); Prov. Čhŏngdžin-si, valley of river Susŏng-čhŏn (110), 21 V 1974, in cow and sheep dung, Z. STEBNICKA (ISEZ); Prov. Kesŏng-si, vicinity of the water-fall Pakjŏn on the Čhŏnma-san (Mt., 13), 6—8 VI 1974, Z. STEBNICKA (ISEZ); Prov. Kangvŏn-do, Kymgang-san (Mts, 53), 18 VI 1974, in cow dung, Z. STEBNICKA (ISEZ).

Remarks. Palaearctic species, rather frequent in D.P.R.K., locally numerous. Collected primarily at light.

Aphodius (Calamosternus) uniplagiatus WATERHOUSE, 1875


(Figs. 85, 86)

Distribution. Taiwan, Korea, Japan — Hokkaido, Honshu (Echigo), Shikoku, Kyushu, Hachijo, Toshima, Kuchinoerabu, Nakanoshima, Takarajima, Iriomote (NAKANE et BABA, 1960; BALTHASAR, 1964; NOMURA, 1966; 1969).

Material examined. 79 ♂♂ and ♀♀. Prov. Kangvŏn-do, Vŏnsan (118), 1 IX 1966, H. SZELĘGIEWICZ and C. DZIADOSZ (ZIW), Kymgang-san (Mts, 53),

16—18 VI 1974, in cow dung, Z. STEBNICKA (ISEZ); Prov. Hvanghe-namdo, vicinity of the water-fall Sujang-san (121) near Hedžu (18), 2 VI 1974, at human excrements, Z. STEBNICKA (ISEZ), valley of river Hakhjön-čhön, 3 VI 1974, in sheep dung, Z. STEBNICKA (ISEZ); Prov. Phjõngan-namdo, distr. Sunan (104), Sõkam-Čõsudži (105), vicinity of the lake, 30 V 1974, in cow dung, Z. STEBNICKA (ISEZ), Nampho (75), Vaudo, 21 VI 1974, Z. STEBNICKA (ISEZ); Prov. Phjõngjang-si, Jongak-san (Mt., 40), 29 V 1974, in sheep dung, Z. STEBNICKA (ISEZ); Prov. Kesõng-si, vicinity of the water-fall Pakjõn on the Čhõnma-san (Mt., 13), 7 VI 1974, in cow dung, Z. STEBNICKA (ISEZ).

Remarks. Common species, in Korea frequent and numerous in various biotopes.



Figs. 96—98. *Aegialia* (*Psammoporus*) *kamtschatica* MOTSCH. 96 — female genitalia, stylus; 97 — aedeagus, lateral view; 98 — paramera, dorsal view. Fig. 99. *A. (Aegialia) comis* (LEWIS) — female genitalia, stylus. Figs. 100—102. Male genitalia, lateral and dorsal view; 100 — *Rhyssemus koreanus* sp. n.; 101 — *Ochodaesus ferrugineus* ESCH.; 102 — *Trox sabulosus* (L.)

Aphodius (Calamosternus) sublimbatus (MOTSCHULSKY, 1860)

Syn.: *Aphodius ussuriensis* PETROVITZ, 1961.

(Figs. 92, 93)

Distribution. East Siberia (Ussuri, Amur), Manchuria, China, Korea, Japan — Hokkaido, Honshu, Shikoku, Kyushu, Tsushima, Kuchinoerabu, Takarajima, Minami-Daitõjima, Toshima, Miyake, Hachijo, Hachijo-Kojima,

Aogashima, Tokara, Amami-Ōshima, Okinawa; Taiwan (Liyutan, Nanshanshi, Liukuei, Lutao Isl., Lanyu Isl.) (BALTHASAR, 1964; NOMURA, 1966; 1969; 1970; 1973).

Material examined. 56 ♂♂ and ♀♀. Japan — Tokyo, Machida, Tsurukawa, 6 VIII 1968, Y. SHIBATA (ISEZ), Yamanashi Pref., near Gozaishi, 27 VII 1967, Y. SHIBATA (ISEZ), Nagano Pref., Kiso Onsen, 1000 m, 19 VI 1970, J. RAZOWSKI (ISEZ); Korea — Prov. Phjŏngan-namdo, distr. Kangsŏ (61), Thesŏng, 24 VIII 1971, at light, J. RAZOWSKI (ISEZ), 13 VI 1974, in cow dung, Z. STEBNICKA (ISEZ), distr. Sunčhŏn (94), Džamo-san, 27 VIII 1971, at light of UV lamps, J. PAWŁOWSKI (ISEZ), distr. Sunan (104), Sŏkam-Čsŭdži (105), 30 V 1974, in cow dung on the banks of the lake, Z. STEBNICKA (ISEZ); Prov. Hvanghe-namdo, valley of river Hakhjŏn-čhŏn near Hedžu (18), 1—3 VI 1974, in cow dung, Z. STEBNICKA (ISEZ), Sinčhŏn (108), 16 IX 1971, in cow dung, J. PAWŁOWSKI (ISEZ); Prov. Čhŏngdžin-si, valley of river Susŏng-čhŏn (110), 21—25 V 1974, in cow and sheep dung, Z. STEBNICKA (ISEZ); Prov. Phjŏngjang-si, Jongak-san (Mt., 40), 29 V 1974, in cow dung on the banks of river, Z. STEBNICKA (ISEZ), 16—20 VIII 1971, at light, J. RAZOWSKI (ISEZ), Phjŏngjang (86), 17 V 1974, at light, A. SZEPTYCKI (ISEZ); Prov. Kangvŏn-do, Kymgang-san (Mts, 53), 16—18 VI 1974, Z. STEBNICKA (ISEZ); Prov. Hamgjŏng-pukto, distr. Kjŏngsŏng (44), valley Onpho-ri (77), 21 VIII 1959, B. PIŚARSKI and J. PRŌSZYŃSKI (ZIW); Prov. Hamgjŏng-namdo, distr. Hjangsan (27), Mhohjangsan (Mts), Hjangam-ri, 18 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. East-Asiatic species inhabiting large area. Forms numerous not distinctly bordered local races, which slightly differ by color and sometimes by shape. Frequent but not numerous in D.P.R.K.

Aegialiini

Aegialia (Aegialia) comis (LEWIS, 1895)

(Fig. 99)

Distribution. USSR — Primorskij Kraj (Suputinka), Japan — Honshu (Nikko), Hokkaido (Sapporo, Kotoni, Bankei, Tokachi, Abashiri); North Korea — Prov. Čhŏngdžin-si, valley of river Susŏng-čhŏn (110) (STEBNICKA, 1977).

Aegialia (Psammoporus) kamtschatica MOTSCHULSKY, 1860

(Figs. 96—98)

Distribution. USSR — Bajkal and Transbajkal Region, Siberia (Daurija), basin of Ussuri river (Birskoje), Kamchatka, Sachalin, Kurile Islands; Japan — Hokkaido, Honshu; North Korea — Prov. Hamgjŏng-pukto, Kvanmo-bong (Mt., 60) (STEBNICKA, 1977).

Material examined. 1 ♀. Prov. Hamgjŏng-pukto, distr. Kjŏngsŏng (44), Mehjang-ri, 4 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

*Psammodiini**Rhyssemus koreanus* sp. n.

(Fig. 100)

Holotype ♂: North Korea, Prov. Hamgjǒng-namdo, distr. Hjangsan (27), Mjohjang-san (Mts), 3 VIII 1959, B. PISARSKI and J. PRÓSZYŃSKI; Paratype ♀, the same data as holotype, in coll. ISEZ.

Description. Length 2.8—3 mm. Elongate-oblong, moderately convex, shining-black; legs, clypeal margin and ventral side reddish. Head strongly convex at middle, edge finely reflexed, clypeus sharply angled each side of broad, moderately deep median emargination, sides weakly arcuate to small genae; front with rounded median convexity, occipital area with two weakly marked carinae and two small tubercles on each side; surface of head everywhere granulated. Pronotum transverse, sides and base margined, setaceous, weakly crenate, marginal setae yellow, short; disc with five transverse, moderately convex carinae and granulated convexity placed on each side near middle of lateral declivity; grooves wide with dense, coarse punctures, surface of carina impunctate. Scutellum small, triangular, alutaceous. Elytra subparallel, humeri sharply dentate; elytral striae moderately deep, intervals slightly convex with a row of moderately large tubercles along outside margin and a row of small tubercles along inner margin. Ventral surface shining; metasternum slightly concave, midline distinct; abdominal segments distinctly delimited by deep, finely and closely punctated transverse grooves; femora with a few punctures bearing yellow hairs. Middle and hind tibiae slender; first segment of posterior tarsus longer than the upper spur and subequal to following three segments combined.

Male. Apical spur of fore tibia bent inward at apex. Aedeagus as in figure 100.

Female. Apical spur of fore tibia straight.

Remarks. *R. koreanus* sp. n. resembles *R. lunatus* PETR. (described from Taiwan) very closely, but these species are separated by differences in the sculpture of pronotum.

*Ochodaeinae**Ochodaeus ferrugineus* (ESCHSCHOLTZ, 1818)

(Fig. 101)

Distribution. Europe, Caucasus, Siberia, Mongolia, Korea (NOMURA, 1942; HORION, 1958; STEBNICKA 1976 et all.).

Material examined. 1 ♂. Prov. Čhǒngdžin-si, Čhǒngdžin (11), 29 VII 1956, M. MAGYAR (HMNH).

*Geotrupinae**Geotrupes (Phelotrupes) auratus* MOTSCHULSKY, 1857

Distribution. USSR — Primorskij Kraj, Kurile Islands (Alechino, Sernovodsk, Kunašir), Japan — Hokkaido, Honshu (Kurokawa, Matsunoyama), Shikoku, Kyushu; Korea (NAKANE et BABA, 1960; MEDVEDEV et ERMOLENKO, 1969; KRYVOLUCKAJA, 1973).

Material examined. 20 ♂♂ and ♀♀. Prov. Hamgjöng-namdo, distr. Hongvön (30), Jönpöng-ri, 8 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), distr. Hjangsan (27), Mjohjang-san (Mts), 24 IX 1966, H. SZELEGIEWICZ and C. DZIADOSZ (ZIW); Prov. Čhöngdžin-si, distr. Purjöng (87), Thomak-töng 20 km N of Čhöngdžin (11), 8 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW); Prov. Phjöngan-namdo, distr. Sunan (104), Junha-ri 40 km W of Phjöng-gjang (86), 13 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), distr. Sunčön (94), Džamo-san and a vicinity of the village Džamo-ri, 27 VIII 1971, J. PAWLOWSKI (ISEZ); Prov. Phjöngjang-si, Jongak-san (Mt., 40), valley of river Sunha-gang, 29 V 1974, under cow dung, Z. STEBNICKA (ISEZ); Prov. Hamgjöng-pukto, locality 20 km NE of Čhöngdžin (11), 5 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Kangvön-do, distr. Vönsan (18), mountain pass Masin-rjöng, 16 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW); Prov. Kesöng-si, vicinity of the water-fall Pakjön on the Čhönma-san (Mt., 13), 7—8 VI 1974, in sheep dung, Z. STEBNICKA (ISEZ), 3 IX 1956, M. MAGYAR (HMNH).

Remarks. Belongs to the group of Manchurian species of rather narrow area of distribution. Fairly common in D.P.R.K.; occurs in the woodland, beetles fly during the day.

Geotrupes (Phelotrupes) laevistriatus MOTSCHULSKY, 1857

Distribution. USSR — Primorskij Kraj, Sachalin, Kurile Islands (Alechino, Goriače lake, Lesnoj river, Lesozavodsk, Kunašir, Sernovodsk, Šikotan, Malokurilskoje), North- and Central China, Manchuria, Korea, Japan — Hokkaido, Honshu (Kurokawa, Mt. Hishigatake, Mt. Futatsumine, Momogawatoge, Mt. Kazakura, Mt. Amakazari), Sado (Suizu, Aoneba-goe, Mt. Donden, Mt. Kinpoku), Shikoku, Kyushu, Okinawa, Yakushima, Iki, Tsushima, Amami-Ōshima, Toshima, Nijima (NAKANE et BABA, 1960; NOMURA, 1966, 1969; MEDVEDEV et ERMOLENKO, 1969; KRYVOLUCKAJA, 1973).

Material examined. 8 ♂♂ and ♀♀. Prov. Janggang-do, distr. Samdžijon (103), Namphote-san (76), 8 IX 1971, in cow dung, J. PAWLOWSKI (ISEZ).

Remarks. An element of Manchurian-fauna. Occurs in the areas with mixed forests in mountainous regions and mountains. Feeds probably in vegetable detritus, sometimes found in cattle excrements. Beetles fly during the day.

*Melolonthinae**Sericini**Gastroserica herzi* (HEYDEN, 1887)

(Figs. 103, 104)

Distribution. China, Manchuria, Taiwan (Hori), Korea — Chosan (125), Usi (116), Kočhang (48), Hverjǒng (25), Kangjasan (49), Andžu (2), Suvǒn (92), Kangnyng (42), Pegam (82), Vǒnsan (118), Pusan (83) (MURAYAMA, 1938; MEDVEDEV, 1952; MURAYAMA, 1954; NAKANE, 1973; NOMURA, 1973).

Material examined. 1 ♂. Prov. Hamgǒng-namdo, distr. Hjangsan (27), Sangvǒn-am (109), 17 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. The occurrence of this species in Japan reported by MURAYAMA (1938, 1954) was not confirmed in the subsequent literature. In the islands of Central Japan occurs very similar species *G. higonía* (LEWIS); the shape of its male copulatory organ is different from this of *G. herzi* (HEYD.) (NOMURA, 1973).

Nipponoserica similis (LEWIS, 1895)

(Figs. 105, 106)

Distribution. Japan — Honshu (Echigo: Sakasamaki), Shikoku (Ehime: Matsuyama), Kyushu (Nagasaki: Nagasaki, Suwanoike; Kumamoto: Nishikoshi), Korea — Čedžu-do (Isl., 3) (MURAYAMA, 1954; NAKANE et BABA, 1960; NOMURA, 1976).

Material examined. 2 ♂♂. Prov. Hvanghe-namdo, vicinity of the water-fall Sujang-san (Mt., 121) near Hedžu (18), 2 VI 1974, under vegetation, Z. STEBNICKA (ISEZ).


Remarks. Species collected in the central part of Peninsula, on the southern slope of the mountain covered with xerophilous vegetation.

Trichoserica polita (GEBLER, 1832)

(Fig. 107, 108)

Distribution. USSR — Bajkal and Transbajkal Region (Čitinskaja Obl.), Bur'atskaja ASSR (Nerčinsk), Primorskij Kraj (Blagoveščensk, Chabarovsk, Askol'd Isl.), Manchuria, Korea — Hverjǒng (25), Sungho (102), Sarivǒn (101), Kangjasan (49) (MURAYAMA, 1938; MEDVEDEV, 1952; MURAYAMA, 1954; NOMURA, 1971).

Material examined. 1 ♂ and 2 ♀♀. Prov. Kesǒng-si, vicinity of the water-fall Pakjǒn on the Čhǒnma-san (Mt., 13), 8 VI 1974, Z. STEBNICKA (ISEZ); Prov.



Figs. 103—115. Male genitalia, lateral and dorsal view; 103 — *Gastroserica herzi* (HEYD.); 104 — *G. higonja* (LEWIS); 105, 106 — *Nipponoserica similis* (LEWIS); 107, 108 — *Trichoserica polita* (GEBL.); 109, 110 — *Sericania fuscolineata* MOTSCH.; 111, 112 — *Maladera (Maladera) gibbiventris* (BRISKE); 113, 114 — *M. (M.) castanea* (ARROW); 115 — *Serica boops* WATERH.

Hamgjöng-namdo, distr. Hjangsan (27), Sangvön-am (109), Mjohjang-san (Mts), 17 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. Specimens were collected on *Rosaceae* bushes.

***Sericania fuscolineata* MOTSCHULSKY, 1860**

(Figs. 109, 110)

Distribution. USSR — Transbaikal Region (Čitinskaja Obl.), Šantarskij Isl., Amur (Nižne-V'atskoje), Daurija, Primorskij Kraj (Askol'd Isl., Vladivostok), North China, Manchuria (four localities recorded in Japanese), Japan — Kyushu (Nagasaki), Ryu-kyu Isl., Tsushima; Korea — Hverjöng (25), Pučhön (river, 91), Kvanmo-bong (Mt., 60) (and 13 not called localities in Japanese) (MURAYAMA, 1938; MEDVEDEV, 1952; MURAYAMA, 1954; NOMURA, 1976).

Material examined. 24 ♂♂ and ♀♀. Prov. Kangvön-do, Kymgang-san (Mts, 53), 17 VI 1974, at light, Z. STEBNICKA (ISEZ), Vönsan (118), 18 VI 1974, Z. STEBNICKA (ISEZ); Prov. Hamgjöng-namdo, distr. Hjangsan (27), Sangvön-am (109), Mjohjang-san (Mts), 17 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), Hjangam-ri, 16—19 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), valley Munsu-tong, 18 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), valley Hapiro, 20—21 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), distr. Hongvön (30), Jönpong-ri, 8 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Hamgjöng-pukto, vicinity of Nadžin (74), 5 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), distr. Kjöngsöng (44), Onpho-ri (77), 3 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Phjöngjang-si, distr. Samsök (106), Tečhön-ri, 22 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), distr. Jongsöng (41), Maram, 29 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Čhön-gdžin-si, Hamgjöng-sanmek (Mts), mountain pass Musan-rjöng, 2 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. Frequent species, locally numerous. In D.P.R.K. found mainly in the mountains. Collected during the day on various plants, comes to light in the evenings.

***Serica boops* WATERHOUSE, 1875**

Syn.: *Ophthalmoserica niijimai* KONTKANEN, 1956.

(Fig. 115)

Distribution. Japan — Honshu (Aomori: Tsuta spa; Fukushima: Kashi, Hutamata; Gunma: Mt. Tanigawa, Hoshi spa, Kiruzumi spa, Odaira; Niigata: Motohashi, Niitsu, Ikenotaira, Shiori-toge, Sakasamaki, Kurokawa, Kanamaru; Nagano: Kuruizawa, Tobira spa, Kiso-Fukushima; Yamanashi: Kiyosato, Ina, Yamanakako, Sagashio spa; Tokyo: Okutama, Mt. Takamizu, Mt. Mitake, Mt. Takao; Kanagawa: Hakone; Shizuoka: Odaru spa, Mt. Fuji; Aichi: Mt. Dando; Gifu: Hirayu; Kyoto: Mt. Hiei; Mie: Misugi; Wakayama: Mt. Koya,

Mt. Ryujin; Hyogo: Maiyasan, Mt. Hyonosen), Shikoku (Ehime: Mt. Nara-bara), Kyushu (Fukuoka: Mt. Wakasugi, Mt. Hikosan, Mt. Shaka), Izu Isl. (Hachijo); Korea — Čedžu-do (Isl., 3) (MURAYAMA, 1938; MEDVEDEV, 1952; MURAYAMA, 1954; NOMURA, 1972).

Material examined. 1 ♂. Prov. Hvanghe-namdo, vicinity of the water-fall Sujang-san (Mt., 121) near Hedžu (18), 2 VI 1974, under stone, Z. STEBNICKA (ISEZ).

Remarks. The occurrence of this species in Manchuria and Primorskij Kraj reported by MURAYAMA (1954) and MEDVEDEV (1952) probably concern a very similar species *S. rosinae* (Pic), whose area of distribution lies more to the north.

Maladera (Maladera) castanea (ARROW, 1913)

Syn.: *Maladera japonica*: MEDVEDEV, 1953 (nec MOTSCHULSKY, 1860).

(Figs. 113, 114)

Distribution. USSR — Primorskij Kraj, Kurile Islands, Caucasus (introduced); North America (introduced); North- and Central China, Japan — Hokkaido (Ishikari), Honshu (Aomori; Akita; Yamagata: Yonezawa; Miyagi: Sendai; Tochigi: Ohtawara; Chiba: Mt. Kiyosumi; Tokyo: Kunitachi, Okutama; Kanagawa: Hiratsuka; Niigata: Kurokawa, Senami, Nakajo, Hirabayashi, Shibata, Kaji, Kama, Yuzawa spa, Maoroshi, Niitsu, Gosen-shi, Gamohara spa, Momozaki-hama, Sasaguchi-hama; Ishikawa: Tsurugi; Nagano: Suwa, Kiso; Shizuoka: Akiba; Gifu; Mie: Misugi; Kyoto; Hyogo: Kobe), Sado (Mt. Kinpoku, Futami, Tassha), Shikoku (Ehime: Sawatani, Ichiu), Kyushu (Fukuoka: Mt. Hikosan, Kokura; Saga; Nagasaki: Kuchinotsu, Tsushima; Kumamoto; Miyazaki: Takanabe; Kagoshima: Kirishima), Ryu-kyu Isl. (Yakushima, Tanegashima), Izu Isl. (Toshima, Nijima, Shikine, Kouzu, Mikura, Miyake, Hachijo, Hachijo-kojima, Aogashima); Taiwan; Korea — Suvŏn (92), Kymgang-san (Mts, 53), Sŏul (93) (MURAYAMA, 1938; NAKANE et BABA, 1960; MURAYAMA, 1954; NOMURA, 1969; 1973; KRYVOLUCKAJA, 1973; MEDVEDEV, 1974).

Material examined. 19 ♂♂ and ♀♀. Prov. Kangvŏn-do, Vŏnsan (118), 1 IX 1966, H. SZEŁĘGIEWICZ and C. DZIADOSZ (ZIW); Prov. Hamgjŏng-namdo, distr. Hjangsan (27), Hjangam-ri, 18 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Phjŏngan-namdo, distr. Sunčhŏn (94), Džamo-san, 27 VIII 1971, J. PAWŁOWSKI (ISEZ), distr. Sunan (104), Sŏkam-Čŏsudži (105), 21 VIII 1971, J. PAWŁOWSKI (ISEZ); Prov. Phjŏngjang-si, Jongak-san (Mt., 40), 20 VIII 1971, J. PAWŁOWSKI (ISEZ), Phjŏngjang (86), 1 VII 1950, BORCHSENIUS (ZIL).

Remarks. MEDVEDEV (1952) considered *M. castanea* as a synonym of *M. japonica* (MOTSCH.), however, they are two different species. The distributional data of *M. castanea* are given above; according to NOMURA (1973), *M. japonica japonica* (MOTSCH.) inhabits Japan exclusively. Both species are enemies of tea and vegetable cultivations and also of orchards; larva gnaws roots of various plants including cultivated ones. One generation per year.

***Maladera (Maladera) gibbiventris* (BRENSKE, 1897)**

(Figs. 111, 112)

Distribution. Central China (Changyang), Taiwan (Wushe, Nanshanchi), Korea — Čungdžu (4), Pujö (79), Čhôngdžu (12), Hedžu (18), Kangnyng (42), Čočhingvön (5), Suvön (92), Söul (93), Kvangjang (43), Kjöngsöng (44), Inčhön (33), Kangge (45) (MURAYAMA, 1938; 1954; NOMURA, 1974).

Material examined. 9 ♂♂ and ♀♀. Prov. Phjôngan-namdo, distr. Kangsö (61), Thesöng, 26 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), distr. Sunan (104), Sökam-Čösudži (105), 30 V 1974, under stones, Z. STEBNICKA (ISEZ); Prov. Phjôngjang-si, distr. Jöngsöng (41), Maram, 26 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), Jongak-san (Mt., 40), 29 V 1974, under stone, Z. STEBNICKA (ISEZ); Prov. Hamgjöng-namdo, distr. Hjangsan (27), Hjangam-ri, 20 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Hvanghe-namdo, Sujang-san (Mt., 121) near Hedžu (18), 2 VI 1974, collected under stones and on the young oak, Z. STEBNICKA (ISEZ); Prov. Kangvön-do, Kymgang-san (Mts, 53) near Kosöng (122), 16 VI 1974, at light, Z. STEBNICKA (ISEZ).

***Maladera (Maladera) formosae* (BRENSKE, 1897)**

(Figs. 116, 117)

Distribution. Taiwan (Wushe, Nanshanchi, Lyutan, Liukuei, Taitsung, Jenai, Tainan, Kenting Park), Korea — Mokpho (66), Suvön (92), Kjöngsöng (44), Söul (93) (MURAYAMA, 1938; 1954; NOMURA, 1974).

Material examined. 6 ♂♂ and ♀♀. Prov. Phjôngan-namdo, distr. Sunan (104), Sökam-Čösudži (105), 21 VIII 1971, at light of UV lamps, J. PAWŁOWSKI (ISEZ); Prov. Kangvön-do, Vönsan (118), 1 V 1966, H. SZEŁĘGIEWICZ and C. DZIADOSZ (ZIW).

***Maladera (Maladera) renardi* (BALLION, 1871)**

Syn.: *Serica spissigrada* BRENSKE, 1897;

Serica motschulskyi BRENSKE, 1897;

Serica nakayamai MURAYAMA, 1938.

(Fig. 118)

Distribution. USSR — Primorskij Kraj (Chabarovsk, Vladivostok), Manchuria, North China, Japan — Honshu (Yamagata: Atsumi spa; Fukushima: Wakamatsu; Ibaragi: Tone; Tokyo: Tamagawa; Kanagawa: Oiso, Kuneguma; Niigata: Shibata, Kurokawa, Sasaguchi-hama, Nakajo, Murakami, Niitsu, Nagaoka; Shimane: Enya), Sado, Shikoku (Tokushima: Kamiryo), Kyushu (Fukuoka: Mt. Hikosan; Nagasaki), Korea — Söul (93), Mudžu (67), Kvesan (62), Kosan (54), Kodže (59), Hongčhön (20), Hedžu (18), Hamhyng (19), Kangge (45), Kildžu (55), Kymčhön (58), Anak (1), Ičhön (33), Kjöngsöng

(44), Purjǒng (87) (MURAYAMA, 1938; MEDVEDEV, 1952; MURAYAMA, 1954; NOMURA, 1973; MEDVEDEV, 1974).

Material examined. 1 ♂ and 1 ♀. Prov. Phjǒngjang-si, Phjǒngjang (86), 26 VIII 1970, at light, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. Beetles are enemies of orchards and cultivations, among others, of broad bean and soia. One generation per year.

***Maladera (Maladera) holosericea* (SCOPOLI, 1772)**

(Figs. 119, 120)

Distribution. Europe, Caucasus, Siberia, Manchuria (numerous localities recorded in Japanese), Korea — Kjǒngsǒng (44), Suvǒn (92), Kesǒng (56), Hamhyng (19), Kangge (45), Kodže (59), Hedžu (18), Hjesan (26), Hverjǒng (25), Kvesan (62), Sǒul (53), Kangnyng (42) (MURAYAMA, 1938; MEDVEDEV, 1952; MURAYAMA, 1954; MEDVEDEV, 1974; STEBNICKA, 1978 et all.).

Material examined. 5 ♂♂ and ♀♀. Prov. Phjǒngjang-si, Phjǒngjang (86), 27 VII 1970, at light, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. Palaearctic species. Beetles are enemies of vegetable cultivations, orchards and field cultivations. Imagines feed on young sprouts, leaves and stamens. One generation per year.

***Maladera (Maladera) cariniceps* (MOSER, 1915)**

(Figs. 121, 122)

Distribution. Manchuria, Japan — Kyushu (Nagasaki), Tsushima; Korea — Mokpho (66), Namvǒn (73), Korjǒng (126), Čhungdžu (4), Hverjǒng (25), Kesǒng (56), Jongdžu (34), Hedžu (18), Čočingvǒn (5), Pujǒ (79), Sǒul (93), Kvesan (62), Inčhǒn (32), Čhǒngdžu (12), Suvǒn (92), Kangnyng (42), Kjǒngsǒng (44), Kvangjang (43), Vǒnsan (118), Čedžu-do (Isl., 3) (MURAYAMA, 1938; 1954; NOMURA, 1973).


Material examined. 6 ♂♂ and ♀♀. Prov. Phjǒngjang-si, Phjǒngjang (86), 17—28 V 1974, under stones on the banks of Tedong-gang (river), Z. STEBNICKA (ISEZ), Jongak-san (Mt., 40) 29 V 1974, valley of river, Z. STEBNICKA (ISEZ).

***Maladera (Maladera) fusania* (MURAYAMA, 1934)**

(Figs. 123, 124)

Distribution. Taiwan (Chihpen), Korea — Mokpho (66), Kvangjang (43), Kunsan (46), Kvangdžu (47), Hedžu (18), Kodže (59), Tongne (111), Mudžu (67), Tegu (112), Čočingvǒn (5), Suvǒn (92), Phohang (81), Sǒul (93), Kangnyng (42), Kjǒngsǒng (44), Hamhyng (19) (MURAYAMA, 1938; 1954; NOMURA, 1974).

Material examined. 16 ♂♂ and ♀♀. Prov. Kangvǒn-do, Vǒnsan (118), 15 VI



Figs. 116—126. Male genitalia, lateral and dorsal view; 116, 117 — *Maladera (Maladera) formosae* (BRŠKE); 118 — *M. (M.) renardi* (BALL.); 119, 120 — *M. (M.) holosericea* (SCOP.); 121, 122 — *M. (M.) cariniceps* (Mos.); 123, 124 — *M. (M.) fusania* (MURAY.); 125, 126 — *M. (M.) orientalis* (MOTSCH).

1974, on the sandy coast of sea, Z. STEBNICKA (ISEZ), 1 IX 1966, H. SZELEGIEWICZ and C. DZIADOSZ (ZIW); Prov. Phjŏngjang-si, Phjŏngjang (86), 14 VI 1974, at light, Z. STEBNICKA (ISEZ); Prov. Kesŏng-si, vicinity of the water-fall Pakjŏn on the Čhŏnma-san (Mt., 13), 7 VI 1974, under stone, Z. STEBNICKA (ISEZ); Prov. Phjŏngan-namdo, distr. Sunan (104), Sŏkam-Čŏsudži (105), vicinity of the lake, 30 V 1974, Z. STEBNICKA (ISEZ); Prov. Hvanghe-namdo, vicinity of the water-fall on the Sujang-san (Mt., 121), 2 VI 1974, under stones, Z. STEBNICKA (ISEZ).

Remarks. Similarly as *M. cariniceps* (Mos.) this species has been collected exclusively in the vicinity of water, most frequently under stones and on the ground, where it spends the day. Active at dusk, comes to light.

Maladera (Maladera) orientalis (MOTSCHULSKY, 1857)Syn.: *Serica salebrosa* BRENSKE, 1897.

(Figs. 125, 126)

Distribution. USSR — Primorskij Kraj, Sachalin; North China (Pekin, Alashan), Mongolia, Manchuria (numerous localities recorded in Japanese), Japan — Hokkaido (Otaru), Honshu (Iwate: Kuji; Miyagi: Sendai; Yamagata: Sendai, Atsumi; Tochigi: Nasu, Ohtawara; Gunma: Yokogawa, Kirizumi; Ibaragi: Mt. Tsukuba; Tokyo: Mt. Takao, Kunitachi, Takamizu, Mt. Kumotori, Tamagawa; Kanagawa: Kamakura, Fujisawa, Tsujido, Hakone; Niigata: Kurokawa, Sasaguchi-hama, Kakuda-hama, Kamiishikawa, Shidai-hama, Mimote, Sakasamaki, Shiunji, Dainichihara, Yoshigahira, Momozaki-hama, Imakawa, Murakami, Senami, Kanamaru, Tsunagi, Yoneyama, Okutadami, Tsuchitaru, Nagaoka, Asagai, Ooyu spa, Agekawa, Shiori-toge; Awa-shima Isl.; Nagano: Mt. Yatsugatake; Shizuoka: Kunoan; Aichi: Nagoya; Gifu: Ishikawa: Tsurugi; Nara: Yamakita; Tokushima: Mt. Tsurugi, Kamiryo), Sado (Umez, Ogura-toge, Mt. Donden), Shikoku (Ehime: Omogo), Kyushu (Fukuoka; Nagasaki), Izu Isl. (Toshima, Nijima); Taiwan; Korea — Hedžu (18) Mokpho (66), Kvandžu (47), Tongne (111), Čongdžu (6), Henam (24), Jōsu (38), Sōul (93), Kangnyng (42), Kosan (54), Kesōng (56), Samčhōnpho (100), Kodže (59), Hamhyng (19), Nadžin (74), Kangge (45), Kvanmo-bong (60), Phohang (81), Mudžu (67), Tegu (112), Čočhingvōn (5), Čhōngdžin (11), Vōndžu (119), Suvōn (92), Jōdžu (39), Inčhōn (32), Kjōngsōng (44), Jōnčhōn (35), Hongčhōn (20), Čedžu-do (Isl., 3) (MURAYAMA, 1938; MEDVEDEV, 1952; MURAYAMA, 1954; NAKANE et BABA, 1960; NOMURA, 1969; 1973; MEDVEDEV, 1974).

Material examined. 34 ♂♂ and ♀♀. Prov. Hamhyng-si, distr. Hamdžu (29), Hyngpong-ri, 12 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), Hyngnam (28), 7—11 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Phjōngnamdo, distr. Sunan (104), Sōkam-Čōsudži (105), 21 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), distr. Sunčhōn (94), Džamo-ri, 27 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), Nampho (75), Usan-ri, 28 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Kangvōn-do, Čhōnne (14), 10 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), Kymgang-san (Mts, 53), 16 VI 1974, Z. STEBNICKA (ISEZ); Prov. Kesōng-si, vicinity of the water-fall Pakjōn on the Čhōnma-san (Mt., 13), 6 VI 1974, Z. STEBNICKA (ISEZ); Prov. Hamgjōngnamdo, distr. Hjangsan (27), Mjohjang-san (Mts), 18 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Hamgjōng-pukto, distr. Kjōngsōng (44), vicinity of Džuyr (17), 25 V 1974, under stones, Z. STEBNICKA (ISEZ); Prov. Phjōngjang-si, Jongak-san (Mt., 40), 24 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), 29 V 1974, Z. STEBNICKA (ISEZ), Tesōng-san (Mts, 124), 21 V 1974, at light, Z. STEBNICKA (ISEZ); Prov. Hvanghe-namdo, Sinčhōn (108), 25 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. This is one of most numerous, widely distributed species of the genus *Maladera* MULS. that occur in Far East. Inhabits mountainous regions. Collected at waters and higher rice fields under stones or directly on the ground; comes to light in the evenings. Beetles are enemies of orchards; imagines destroy young sprouts and leaves, larvae — roots of various trees and shrubs. One generation per year.

***Maladera (Maladera) schonfeldti* (MURAYAMA, 1937)**

(Figs. 127, 128)

Distribution. Korea — Mokpho (66), Suvŏn (92), Korjŏng (126), Kesŏng (56), Kangsŏ (61), Hamhyng (19) (MURAYAMA, 1938; 1954).

Material examined. 45 ♂♂ and ♀♀. Prov. Čhŏngdžin-si, Čhŏngdžin (11), 21 V 1974, at light, Z. STEBNICKA (ISEZ); Prov. Hamhyng-si, Hyngnam (28), 7 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Phjŏngjang-si, Phjŏngjang (86), 20 V 1974, at light, Z. STEBNICKA (ISEZ).

Remarks. This species has been known up to now only from Korea. Local, relatively numerous, collected at light. Body surface of the beetles brown, brownish-black or black, covered by a characteristic grey coating. Description of this species is not contained in the present paper since the representatives of the genus *Maladera* MULS. are very similar externally and indicate a great individual variability; the difference in the structure of their male copulatory apparatuses is the only criterium on which they are properly determinable.

***Maladera (Maladera) okamotoi* (MURAYAMA, 1938)**

(Figs. 129, 130)

Distribution. China, Manchuria (two localities recorded in Japanese), Korea — Namvŏn (73), Jongdžu (34), Čhŏngdžu (12), Čhungdžu (4), Čŏchingvŏn (5), Suvŏn (92) (MURAYAMA, 1938; 1954).

Material examined. 5 ♂♂ and ♀♀. Prov. Janggang-do, distr. Hjesan (26), vicinity of the village Dŏngha-ri, 31 VIII 1971, at light of UV lamps, J. RAZOWSKI (ISEZ); Prov. Phjŏngjang-namdo, distr. Sunan (104), Junha-ri, 3 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW).


Remarks. This species was collected only at light.

***Maladera (Eumaladera) nitidiceps* NOMURA, 1967**

(Figs. 131, 132)

Distribution. Japan — Ryu-kyu (Iriomote-jima) (NOMURA, 1973).

Material examined. 2 ♂♂ and 1 ♀. Prov. Phjŏngjang-si, Phjŏngjang (86), 17 IX 1959, B. PISARSKI and J. PRÓSZYŃSKI (ZIW); Prov. Hamgjŏng-namdo, distr. Hjangsan (27), Hjangam-ri, 20 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).



Figs. 127—134. Male genitalia, lateral and dorsal view; 127, 128 — *Maladera* (*Maladera*) *schonfeldti* (MURAY.); 129, 130 — *M. (M.) okamotoi* (MURAY.); 131, 132 — *M. (Eumaladera)* *nitidiceps* NOM.; 133, 134 — *Hoplia* (*Euchromoplia*) *aureola* (PALL.). Figs. 135, 136. *H. (Decamera)* *djukini* JACOB. 135 — last segment and claw of posterior tarsus; 136 — antenna. Figs. 137, 138. *Hilyotrogus bicolorus* (HEYD.) — male genitalia, lateral and dorsal view. Figs. 139, 140. *Ectinohoplia rufipes* (MOTSCH.); 139 — last segment and claw of posterior tarsus; 140 — inner claw of anterior tarsus. Figs. 141, 142. *Hoplosternus incanus* MOTSCH.; 141 — prosternal process; 142 — pigidium

Remarks. Subgenus *Eumaladera* NOM. contains two Japanese species inhabiting Ryu-kyu Archipelago: *M. (E.) nitididorsis* NOM. and *M. (E.) nitidiceps* NOM. These species differ from the representatives of the subgenus *Maladera* s. str. by following features: hind tibia broad and flat with longitudinal serrated ridge from base to near apex on outer side. Hind tarsi with short setae on ventral side. Anterior margin of hind femur finely serrate. Dorsal surface partially shining. Color red-brown to dark red-brown. *M. (E.) nitidiceps* is a new species to the Korean fauna.

Hopliini

Hoplia (Euchromoplia) aureola (PALLAS, 1781)

(Figs. 133, 134)

Distribution. East Siberia, Mongolia, Manchuria, North China, Korea (MEDVEDEV, 1952; 1974; 1976; ENDRÖDI, 1965; 1967; 1969; 1971).

Material examined. 66 ♂♂ and ♀♀. Prov. Kesöng-si, vicinity of the waterfall Pakjön on the Čhönma-san (Mt., 13), 6 VI 1974, Z. STEBNICKA (ISEZ); Prov. Phjongan-namdo, distr. Sunčhön (94), Džamo-ri, 27 V 1965, N. MROCZKOWSKI and A. RIEDEL (ZIW), distr. Sunan (104), Sökam-Čösudži (105), 21 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Phjongjang-si, distr. Jongsöng (41), Maram, 29 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Hamgjöng-namdo, distr. Hongvön (30), Jönpong-ri, 8 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. An element of Manchurian fauna. Beetles appear in great numbers, cover various herbaceous plants and bush scrubs; they are enemies of cabbage and rice cultivations. In D.P.R.K. collected frequently in the vicinity of plough-lands, at stream borders, on bushes and flowers.

Hoplia (Decamera) djukini JACOBSON, 1914

(Figs. 135, 136)

Distribution. USSR — Primorskij Kraj (Vorošilov-Ussurijskij, Šmakovka, vicinity of Spassk, Vinogradovka, Tiutiche river, Tumen-ula river), (MEDVEDEV, 1952).

Material examined. 1 ♀. Prov. Hamhyng-si, Hyngnam (28), 11 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. According to MEDVEDEV (1952) beetles occur on the banks of inland waters and at sea shore, often locally in Primorskij Kraj. Species new to the Korean fauna.

Ectinohoplia rufipes (MOTSCHULSKY, 1860)

(Figs. 139, 140)

Distribution. USSR — Amur, Primorskij Kraj, Sachalin, Kurile Islands (Sernovodsk, Alechino, Goriače Lake, Malokurilsk, Veslovskij Pen.) North- and East China, Korea, Japan — Hokkaido, Honshu, Shikoku, Kyushu, (MEDVEDEV et ERMOLENKO, 1969; KRYVOLUCKAJA, 1973; MEDVEDEV, 1974).

Material examined. 2 ♀♀. Prov. Kesōng-si, vicinity of the water-fall Pakjōn on the Čhōnma-san (Mt., 13), 8 VI 1974, Z. STEBNICKA (ISEZ).

Remarks. Beetles inhabit wooded areas, occur in great numbers in Primorskij Kraj. By eating leaves of apple-trees, plum-trees and other fruit-trees often cause great damages in orchards.

*Melolonthini**Hoplosternus incanus* MOTSCHULSKY, 1853

(Figs. 141, 142)

Distribution. USSR — Primorskij Kraj (Vorošilov-Ussurijskij, Vladivostok, Tumen-ula river), North China (to Peking), Manchuria (twelve localities recorded in Japanese), Japan — Kyushu; Korea — Mokpho (66), Kangjasan (49), Suvōn (92), Sičhang (97), Čočhingvōn (5), Kjōngsōng (44), Kymgang-san (Mts, 53), Ičhōn (33), Hedžu (18), Hamhyng (19), Musan (70), Kangge (45), Sōnčhōn (95), Sinyidžu (96), Purjōng (87) (and 13 not called localities recorded in Japanese) (MEDVEDEV, 1951; MURAYAMA, 1954).

Material examined. 4 ♂♂ and ♀♀. Prov. Phjōngjang-si, Jongak-san (Mt., 40), 20 VIII 1971, at light of UV lamps, J. RAZOWSKI (ISEZ); Prov. Hamgjōng-punkto, distr. Kjōngsōng (44), Onpho-ri (77), 14—15 VIII 1959, B. PIŚARSKI and J. PRÓSZYŃSKI (ZIW).

Remarks. Similar to the majority of representatives of the tribe *Melolonthini*, this species appears exclusively in the evenings and at the beginning of nights. Linked to woodland, collected only at light.

*Heptophyllini**Hilyotrogus bicolorus* (HEYDEN, 1887)

(Figs. 137, 138)

Distribution. USSR — Primorskij Kraj (Vladivostok, Olga), Manchuria, Korea — Kangjasan (49), Korjōng (126), Kymgang-san (Mts, 53) (and 13 not called localities recorded in Japanese) (MEDVEDEV, 1951; MURAYAMA, 1954).

Material examined. 1 ♂ and 2 ♀♀. Prov. Phjōnggan-namdo, distr. Sunčhōn (94), Džamo-san, 27 VIII 1971, at light, J. PAWŁOWSKI and A. SZEPTYCKI (ISEZ).

Remarks. Rare species, probably mountainous, collected as isolated individuals. The not-deciphered names of the localities in Korea given by MURAYAMA (1954) in the Japanese language, mainly indicate mountains or eminences (—„san” ending). The names of particular summits were created according to designations by local inhabitants, therefore the identification of them is very difficult.

Rhizotrogini

Apogonia cupreoviridis KOLBE, 1886

(Figs. 143, 144)

Distribution. Japan, Manchuria (seven localities recorded in Japanese), Korea — Vando (120), Kvangjang (43), Sunčhön (94), Tongne (111), Kvangdžu (47). Mokpho (66), Namvön (73), Kangjasan (49), Mudžu (67), Čongdžu (6), Ponghva-ri (88), Uldžin (117), Čočingvön (5), Thosan (114), Suvön (92), Kjöngsöng (44). Pisylsan (89), Čunchön (8), Vöndžu (119), Hongčhön (20), Ičhön (33), Vönsan (118), Hedžu (18), Kečhön (57), Čongphjöng (10), Jöngvön (37) (and nine not called localities recorded in Japanese) (MURAYAMA, 1954).

Material examined. 82 ♂♂ and ♀♀. Prov. Hvanghe-namdo, Sujang-san (Mt., 121), vicinity of the water-fall, 3 VI 1974, Z. STEBNICKA (ISEZ), Sinčhön (108), 26 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Phjöngan-namdo, Nampho (75), Vaudo, 21 VI 1974, Z. STEBNICKA (ISEZ), 28 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), Usan-ri, 28 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), distr. Suhan (104), Sökam-Čösud (105), vicinity of the lake, 30 V 1974, Z. STEBNICKA (ISEZ), distr. Kangsö (61), Thesöng, 26 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Kangvön-do, Vönsan (118), 15 VI 1974, at light, Z. STEBNICKA (ISEZ), Kymgang-san (Mts, 53) near Kosöng (122), 16—18 VI 1974, Z. STEBNICKA (ISEZ); Prov. Phjöngjang-si, Tesöng-san (Mts, 124), 20 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), 16 V 1974, Z. STEBNICKA (ISEZ), Phjöngjang (86), 26 VI 1974, Z. STEBNICKA (ISEZ), Jongak-san (Mt., 40), 29 V 1974, under stones, Z. STEBNICKA (ISEZ), 24 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), distr. Samsök (106), Söngmun-ri, 22 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Janggang-do, distr. Hjesan (26), 11 IX 1971, at light of UV lamps, J. RAZOWSKI (ISEZ).


Remarks. Species frequent and numerous in D.P.R.K. Collected mainly under stones, on plants or directly on the ground, also at light. Most frequently occurs at sea-shore, in the valleys of mountainous rivers and streams and in the vicinity of lakes.

Apogonia cribricollis BURMEISTER, 1855

(Figs. 145, 146)

Distribution. Manchuria (six localities recorded in Japanese), Korea — Suvön (92), Kjöngsöng (44) (MURAYAMA, 1954).

Material examined. 6 ♂♂ and ♀♀. Prov. Hamgjöng-namdo, distr. Hjangsan



Figs. 143—156. Male genitalia, lateral and dorsal view; 143, 144 — *Apogonia cupreoviridis* KOLBE; 145, 146 — *A. cribricollis* BURM.; 147, 148 — *Lasiopsis manchuricus* MURAY.; 149, 150 — *Brahmina rubetra* (FALD.); 151, 152 — *Sophrops heydeni* (BRISKE); 153, 154 — *S. striata* (BRISKE); 155, 156 — *Miridiba koreana* NIJ. et KIN.

(27), Hjangam-ri, Mjohjang-san (Mts), 19 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Phjôngjang-si, Phjôngjang (86), town park Moran-bong, 29 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. Species much more rare than the former one; nothing is known of the ecology of its. Differs from *A. cupreoviridis* KOLBE by smaller body size and punctation of elytra.

***Brahmina rubetra* (FALDERMANN, 1835)**

(Figs. 149, 150)

Distribution. North China (to Peking), Manchuria (MEDVEDEV, 1951; MURAYAMA, 1954).

Material examined. 1 ♂ and 1 ♀. Prov. Phjôngjang-si, Phjôngjang (86), town park Moran-bong, 27 VII 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW); Prov. Hamgjông-namdo, distr. Hjangsan (27), Mjohjang-san (Mts), 9 VIII 1959, B. PISARSKI and J. PRÓSZYŃSKI (ZIW).

Remarks. A new species to the Korean fauna. It is probably an element of the Manchurian fauna of a narrow range.

***Lasiopsis manchuricus* MURAYAMA, 1941**

(Figs. 147, 148)

Distribution. Manchuria (two localities recorded in Japanese) (MURAYAMA, 1954).

Material examined. 1 ♂. Prov. Phjônggan-namdo, distr. Sunan (104), Junhari, 13 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW).

Remarks. Species new to the Korean fauna, up to now known only from Manchuria. It differs from the other Far-Eastern species of the genus *Lasiopsis* ER. by the structure of male copulatory apparatus.

***Sophrops heydeni* (BRENSKE, 1892)**

(Figs. 151, 152)

Distribution. USSR — Primorskij Kraj (Černigovka, Jakovlevka, Vorosilov-Ussurijskij, Tumen-ula river), Korea — Mokpho (66), Peksan (84), Kangjasan (49), Tegu (112), Mudžu (67), Čočhingvön (5), Suvön (92), Kjôngsöng (44), Inčhön (32), Hongčhön (20) (and nine not called localities recorded in Japanese) (MEDVEDEV, 1951; MURAYAMA, 1954).

Material examined. 1 ♂. Prov. Phjônggan-namdo, distr. Sunan (104), Sökam-Čösudži (105), vicinity of the lake, 27 VIII 1971, at light of UV lamps, J. PAWŁOWSKI (ISEZ).

Remarks. Beetles active in the evenings.

Sophrops striata (BRENSKE, 1892)

(Figs. 153, 154)

Distribution. China, Korea — Kangjasan (49), Korjǒng (126), Suvǒn (92), Hvado (127), Sokǒho (129), Kjǒngsǒng (44), Sarivǒn (101), Sinsang (128), Inǒhǒn (32), Andžu (2), Sǒnǒhǒn (95), Sǒnsan (130) (and six not called localities recorded in Japanese) (MEDVEDEV, 1951; MURAYAMA, 1954).

Material examined. 6 ♂♂ and ♀♀. Prov. Kangvǒn-do, Kymgang-san (Mts, 53), 16 VI 1974, at light, Z. STEBNICKA and A. SZEPTYCKI (ISEZ); Prov. Phjǒnggan-namdo, distr. Kangsǒ (61), Thesǒng, vicinity of the lake, 13 VI 1974, under stone, Z. STEBNICKA (ISEZ); Prov. Hvanghe-namdo, vicinity of the water-fall Sujang-san (Mt., 121) near Hedžu (18), 2 VI 1974, under stone, Z. STEBNICKA (ISEZ).

Remarks. Species nocturnal, beginning activity soon after sunset. During the day the beetles stay under stones and plants, come to light in the evenings.

Miridiba koreana NILJIMA et KINOSHITA, 1923

(Figs. 155, 156)

Distribution. Manchuria (five localities recorded in Japanese), Korea — Mokpho (66), Jongdžu (34), Čungdžu (4), Čočhingvǒn (5), Suvǒn (92), Kjǒngsǒng (44), Vǒnsan (118), Sepho (98), Andžu (2), Kangge (45), Sinphjǒng (99) (MURAYAMA, 1954).

Material examined. 14 ♂♂ and ♀♀. Prov. Phjǒngjang-si, Phjǒngjang (86), town park Moran-bong, 14 VI 1974, at light, Z. STEBNICKA (ISEZ).

Remarks. This is one of three known species of the genus *Miridiba* REITT. inhabiting North China, Manchuria and Korea. Beetles are active at dusk and in the nights, collected only at light (cover the shades of street lamps).

Holotrichia diomphalia (BATES, 1888)

(Fig. 157)

Distribution. USSR — Transbaikal Region (Čitinskaja Obl.), Amur, Ussuri, Primorskij Kraj (Tarasovka), Sachalin; North China (to Peking), Manchuria (24 localities recorded in Japanese), Japan, Korea — Mokpho (66), Vando (120), Kvangjang (43), Sunǒhǒn (94), Tongne (111), Peksan (84), Tegu (112), Kangjasan (49), Kongdžu (131), Kymgang-san (Mts, 53), Kvangdžu (47), Čǒngdžu (6), Suvǒn (92), Kjǒngsǒng (44), Kosan (54), Hongǒhǒn (20), Čunǒhǒn (8), Jǒnǒhǒn (35), Iǒhǒn (33), Kangnyng (42), Kočchang (48), Jongdǒk (36), Vǒndžu (119), Koksan (64), Čǒngdžu (12), Hamhyng (19), Phjǒngjang (86), Jǒngvǒn (37), Sǒnǒhǒn (95), Kangge (45), Mjǒngǒhǒn (68), Čǒngsǒng (9), Kildžu (55) (MEDVEDEV, 1951; MURAYAMA, 1954; MEDVEDEV, 1974; KALININA, 1977).

Material examined. 40 ♂♂ and ♀♀. Prov. Kangvön-do Kymgang-san (Mts, 53), 16 VI 1974, at light, Z. STEBNICKA (ISEZ); Prov. Hamgjöng-namdo, distr. Hjangsan (27), Hjangam-ri, Mjohjang-san (Mts), 21 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), valley Munsu-tong, 18 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), 3 VIII 1959, B. PISARSKI and J. PRÓSZYŃSKI (ZIW); Prov. Phjööngjang-si, distr. Samsök (106), Söngmun-ri, 22 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), Tečhön-ri, 22 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), Jongak-san (Mt., 40), 24 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Čhöngdžin-si, mountain chain Hamgjöng-sanmek, pass Musan-rjöng, 2 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Hamhyng-si, Hamhyng (19), 7 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); South Korea, Čedžu-do (Isl., 3), coll. LGOCKI (ISEZ).

Remarks. Numerous species of the genus *Holotrichia* HOPE, which mainly inhabit East Asia are very similar externally and difficult to determine, however, the morphology of male copulatory apparatuses of particular species is distinctly differentiated. Bionomy of the representatives of this genus is very little known. Beetles are active at evenings and nights, however, nearly exclusively females were collected by means of electric light. This indicates that mainly females fly. Imagines probably feed on leaves and young tree-sprouts. The developmental cycle of *H. diomphalia* (BAT.) takes two years on an average; larva winters twice. Larvae destroy the cultivations of rice, wheat, barley, millet, maize, pulse plants, flax, hemp, sugar cane and potatoes.


Holotrichia ernesti REITTER, 1902

(Fig. 158)

Distribution. USSR — Amur, Ussuri, Primorskij Kraj (Chanka Lake), North China, Mongolia, Manchuria (Mukden, Harbin), Korea — Namvön (73), Korjöng (126), Suvön (92), Kjööngsöng (44) (MEDVEDEV, 1951; MURAYAMA, 1954; KALININA, 1977).

Material examined. 6 ♂♂ and ♀♀. Prov. Hamgjöng-punkto, distr. Kjööngsöng (44), Džuyr (17), Jonghen, 25 V 1974, Z. STEBNICKA (ISEZ); Prov. Čhöngdžin-si, Purjöng (87), valley of river, 22 V 1974, under stones, Z. STEBNICKA (ISEZ); Prov. Hvanghe-namdo, Sujang-san (Mt., 121) near Hedžu (18), 2 VI 1974, under stone, Z. STEBNICKA (ISEZ); Prov. Hamgjöng-namdo, distr. Hongvön (30), Jönpong-ri, 8 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Hamhyng-si, distr. Hamdžu (29), Hyngpong-ri, 12 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Phjööngjang-si, Phjööngjang (86), 24 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. Fairly frequent but not numerous species, beginning activity soon after sunset; during the day the beetles stay in the ground or under stones, come to light in the evenings. Probably linked to watersides.



Figs. 157—164. Male genitalia, lateral and dorsal view; 157 — *Holotrichia diomphalia* (BAT.); 158 — *H. ernesti* REITT.; 159, 160 — *Eotrichia titanis* (REITT.); 161, 162 — *Holotrichia parallella* (MOTSCH.); 163, 164 — *H. picea* WATERH.

***Holotrichia parallella* (MOTSCHULSKY, 1854)**

Syn.: *Holotrichia morosa* WATERHOUSE, 1875.

(Figs. 161, 162)

Distribution. USSR — Sachalin, Primorskij Kraj (Tumen-ula), Japan — Honshu (Kurokawa), Tsushima; North- and Central China, Tibet, Manchuria (six localities recorded in Japanese), Korea — Čedžu-do (Isl. 3), Kvangjang (43), Vando (120), Kvangdžu (47), Sunčhŏn (94), Mokpho (66), Tongne (111), Kangjasan (49), Namvŏn (73), Pungan (85), Kangdžin (50), Hadong (21), Kjŏngdžu (51), Čhŏngdžu (12), Tegu (112). Čhŏngan (7), Hvengjang (22), Tedžŏn

(113), Suvŏn (92), Jongdžu (34), Kimpho (52), Inčhŏn (32), Kjŏngsŏng (44), Korjŏng (126), Čočhingvŏn (5), Čhunčhŏn (8), Kymgang-san (Mts, 53), Hvačhŏn (23) (MEDVEDEV, 1951; NAKANE et BABA, 1960; MURAYAMA, 1954; KALININA, 1977).

Material examined. 6 ♂♂ and ♀♀. Prov. Kangvŏn-do, Kymgang-san (Mts, 53), 17 VI 1974, at light, Z. STEBNICKA and A. SZEPTYCKI (ISEZ); Prov. Phjŏngjang-si, Jongak-san (Mt., 40), 18 VIII 1971, at light of UV lamps, J. RAZOWSKI (ISEZ), Phjŏngjang (86), 21 VII 1959, B. PISARSKI and J. PRÓSZYŃSKI (ZIW).

Holotrichia picea WATERHOUSE, 1875

Syn.: *Holotrichia infantula* MEDVEDEV, 1951.

(Figs. 163—166, 172)

Distribution. USSR — Promorskij Kraj (Chasan, Sidemi, Barabaš, Tumen-ula, Razdolnoj river), Japan — Hokkaido, Honshu (Momozaki-hama, Mt. Yakemine, Kyoto), Awa-shima, Kyushu; Manchuria, Korea — Kangjasan (49), Čedžu-do (Isl., 3) (MEDVEDEV, 1951; NAKANE et BABA, 1960; MURAYAMA, 1954; KALININA, 1977).

Material examined. 10 ♀♀. Prov. Phjŏngjang-si, distr. Samsŏk (106), Tečhŏn-ri, 22 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), Jongak-san (Mt., 40), 24 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Hamgjŏng-namdo, distr. Hjangsan (27), Mjohjang-san (Mts), valley Manphok-tong, 19 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Phjŏngan-namdo, distr. Sunan (104), Sŏkam-Čŏsudži (105), 21 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. Species spread in the northern part of the Peninsula, not numerous. Collected at light.

Holotrichia oblita (FALDERMANN, 1835)


Syn.: *Holotrichia amplipennis* FAIRMAIRE, 1887.

(Figs. 167, 168)

Distribution. Manchuria, Japan — Honshu (MEDVEDEV, 1951).

Material examined. 12 ♀♀. Prov. Hamgjŏng-namdo, distr. Hjangsan (27), Hjangam-ri, Mjohjang-san (Mts), 29 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Phjŏngjang-si, Phjŏngjang (86), 12—22 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), Jongak-san (Mt., 40), 24 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Phjŏngan-namdo, distr. Kangsŏ (61), Thesŏng, 26 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Kesŏng-si, vicinity of the water-fall Pakjŏn on the Čhŏnma-san (Mt., 13), 16 VI 1974, under stones, Z. STEBNICKA (ISEZ); Prov. Kangvŏn-do, Kymgang-san (Mts, 53), 17 VI 1974, at light, Z. STEBNICKA and A. SZEPTYCKI (ISEZ), Samil-pho (123), vicinity of the lake, 19 VI 1974, J. PAWŁOWSKI and A. SZEPTYCKI (ISEZ); Prov. Hvanphe-namdo, Sinčhŏn (108), 25 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. This species reported by MEDVEDEV (1951) from Primorskiy Kraj, according to KALININA (1973) does not occur on the territory of USSR. New to the Korean fauna, widely distributed in northern Provinces. Fairly frequent, unfortunately only females were collected. Probably linked to water-sides. Evening-active, comes to light, during the day the beetles hide in the ground or under stones.



Figs. 165, 166. *Holotrichia picea* WATERH.; 165 — female pigidium; 166 — apex of posterior tibia. Figs. 167, 168. *H. oblita* (FALD.); 167 — female pigidium; 168 — apex of posterior tibia. Figs. 169—171. *H. inelegans* (LEWIS); 169 — female pigidium; 170 — apex of posterior tibia; 171 — last segment of maxillary palpus. Fig. 172. *H. picea* WATERH. — last segment of maxillary palpus

Holotrichia inelegans (LEWIS, 1895)

(Figs. 169—171)

Distribution. USSR — Primorskiy Kraj (Tatjanovka, Kamen-Rybolov, Troickoje, Jakovlevka, Ussurijsk, vicinity of Chabarovsk, Nižn. Tambovskoje), Sachalin; Manchuria (Mukden), Japan — Hokkaido, Honshu, Kyushu (MEDVEDEV, 1951; KALININA, 1977).

Material examined. 1 ♀. Prov. Čhôngdžin-si, Čhôngdžin (11), 21 V 1974, at light, Z. STEBNICKA (ISEZ).

Remarks. This species has been probably reported from Korea by MURAYAMA (1954) as *H. kiotoensis* BRISKE, however, due to lack of the material I can not check this report. A detailed revision of the genus *Holotrichia* HOPE is desirable, with reference to the materials from China and to the reports done previously by the Chinese (CHANG YOU-WEI, 1965) and Japanese authors.

Eotrichia titanis (REITTER, 1902)

(Figs. 159, 160)

Distribution. USSR — Primorskij Kraj (Černigovka, Kamen-Rybolov, Novo-Nikolskoje, Vorosilov-Ussurijskij, Ula-che and Tumen-ula rivers), Manchuria, Korea — Tegu (112), Suvŏn (92), Vando (120), Kjŏngsŏng (44), Vŏndžu (119), Kesŏng (56), Čhunčhŏn (8), Čhôngdžu (12), Mengsan (69), Hamhyng (19), Sinyidžu (96), (MEDVEDEV, 1951; MURAYAMA, 1954; MEDVEDEV, 1974).

Material examined. 1 ♂ and 4 ♀♀. Prov. Phjŏngan-namdo, distr. Sunan (104), Sŏkam-Čŏsudži (105), 21 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), distr. Kangsŏ (61), Thesŏng, 26 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Phjŏngjang-si, Tesŏng-san (Mts, 124), 20 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. Bionomics similar to that of *H. diomphalia* (BAT.). Beetles are evening-active, come to light. Larvae are enemies of cultivations.

Rutelinae

Rutelini


Popillia indigonacea MOTSCHULSKÝ, 1853

(Figs. 173, 174)

Distribution. USSR — Amur, Ussuri, Bureinskij Chrebet, Primorskij Kraj (Sedanka, Černigovka), Manchuria, East China, Korea, Japan — Ryu-kyu Isl.; Taiwan (MEDVEDEV, 1949; MACHATSCHKE, 1972; MEDVEDEV, 1974).

Material examined. 6 ♂♂ and ♀♀. Prov. Phjŏngan-namdo, distr. Sunan (104), Sŏkam-Čŏsudži (105), 2 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW); Prov. Hamgjŏng-pukto, Kjŏngsŏng (44), 11 IX 1966, H. SZELEGIEWICZ and C. DZIADOSZ (ZIW); Prov. Hamgjŏng-namdo, Pučhŏn (91), 2 VIII 1950 (ZIL); Prov. Janggang-do, distr. Hjesan (26), vicinity of the village Dŏng-ha-ri, 2 IX 1971, J. PAWŁOWSKI (ISEZ); Prov. Kangvŏn-do, Kymgang-san (Mts, 53), 18 VI 1974, Z. STEBNICKA (ISEZ).

Remarks. Species widely spread but not numerous in D.P.R.K. Collected with *P. atrocoerulea* BAT. mainly in the flowers of *Rosa* L. According to MEDVEDEV (1974) this species is an enemy of cotton in China and Taiwan.



Figs. 173—182. Male genitalia, lateral and dorsal view; 173, 174 — *Popillia indigonacea* MOTSCH.; 175, 176 — *P. atrocoerulea* BAT.; 177, 178 — *P. japonica* NEWM.; 179, 180 — *P. quadriguttata* (FABR.); 181, 182 — *P. ruficollis* KR.

***Popillia atrocoerulea* BATES, 1888**

(Figs. 175, 176)

Distribution. USSR — Primorskij Kraj (Chasan, Tumen-ula), North- and Central China, Taiwan, Korea — Vönsan (118) (MEDVEDEV, 1949; MACHATSCHKE, 1972; MEDVEDEV, 1974).

Material examined. 43 ♂♂ and ♀♀. Prov. Phjôngjang-si, Tesông-san (Mts, 124), 26 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Hamgjông-namdo, distr. Hjangsan (27), Mjohjang-san (Mts), 22 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), Hjangam-ri, 21 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Kangvôn-do, Kymgang-san (Mts, 53), 16—18 VI 1974, Z. STEBNICKA (ISEZ), Čhônne (14), 10 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. Similarly to the former one this species inhabits areas with abundant vegetation. It flies in the daytime, feeds on flowers, leaves and also on fruits of numerous plants. More numerous in southern Provinces of D.P.R.K. Collected on flowers of *Rosa* L. One-colored typical form occurs with the sporadically occurring ab. *biplagiata* KR.

Popillia japonica NEWMANN, 1838

(Figs. 177, 178)

Distribution. USSR — Kurile Islands (Kunašir), North China, Japan — Hokkaido, Honshu (Ginzandaira, Yoshigahira, Nanatani), Awa-shima, Sado (Sukunegi, Umezu, Mt. Donden), Shikoku, Kyushu; North America (New Jersey) and Canada (introduced) (MEDVEDEV, 1949; NAKANE et BABA, 1960; MACHATSCHKE, 1972; KRYVOLUCKAJA, 1973; MEDVEDEV, 1974).

Material examined. 17 ♂♂ and ♀♀. Prov. Čhôngdžin-si, Purjông (87), valley of river, 22 V 1974, Z. STEBNICKA (ISEZ); Prov. Phjôngjang-si, Phjôngjang (86), 21 VII 1959, B. PISARSKI and J. PRÓSZYŃSKI (ZIW).

Remarks. According to MEDVEDEV (1949) *P. japonica* is an endemic, Japanese species introduced to China and to the American Continent, where it considerably multiplied and caused great damages in cultures. Nevertheless in Japan it never does occur in great masses, consequently it does not have an economic significance. Beetles are enemies of gardens and fruit-trees, larvae damage roots of various plants. This species was not reported from Korea up the present. Often mistaken as *P. quadriguttata* (FABR.) because seems not different externally.

Popillia quadriguttata (FABRICIUS, 1787)

(Figs. 179, 180)

Distribution. USSR — Primorskij Kraj (Chabarovsk), Manchuria (Harbin, Mukden), North China (Peking, Tianjin), Central China (Kansu), Korea, Taiwan, Vietnam (MEDVEDEV, 1949; MACHATSCHKE, 1972; MEDVEDEV, 1974).

Material examined. 5 ♂♂ and ♀♀. Prov. Hamgjông-namdo, distr. Hongvôn (30), Jônpong-ri, 8 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Phjôngjang-si, Phjôngjang (86), 21 VII 1959, B. PISARSKI and J. PRÓSZYŃSKI (ZIW).

Remarks. This species occurs mainly in the river-valleys. Beetles are enemies of vegetable cultures (pulse plants, cabbage, potatoes), fruit-trees, rye, wheat and vineyards.

***Popillia ruficollis* KRAATZ, 1892**

(Figs. 181, 182)

Distribution. South Manchuria, East China (to Shanghai) (MEDVEDEV, 1949; MACHATSCHKE, 1972).

Material examined. 1 ♂. Prov. Phjǽngjang-si, Phjǽngjang (86), 29 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. Species very similar to the former one. Described by KRAATZ (1892) as *P. quadriguttata* var. *ruficollis*, formed as separate species by MEDVEDEV (1949). New to the Korean fauna.

***Phyllopertha horticola* (LINNAEUS, 1758)**

(Figs. 183, 184)

Distribution. Europe, Caucasus, Siberia, Mongolia, Manchuria, North China (MEDVEDEV, 1949; MACHATSCHKE, 1972; MEDVEDEV, 1974; STEBNICKA, 1978 et all.).

Material examined. 6 ♂♂ and ♀♀. Prov. Phjǽgan-namdo, distr. Sunan (104), Sǽkam-Čǽsudži (105), 21 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. Palaearctic species, not recorded from Korea up to the present. Beetles, sometimes very numerous feed on various plants; they are enemies of gardens and orchards.

***Spilota plagicollis* FAIRMAIRE, 1886**

(Figs. 185, 186)

Distribution. USSR — Primorskij Kraj (Vladivostok), North- and Central China, Korea (MEDVEDEV, 1949; MACHATSCHKE, 1972).

Material examined. 16 ♂♂ and ♀♀. Prov. Hamgjǽng-namdo, distr. Hjangsan (27), Hjangam-ri, 18 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), Mjohjangsan (Mts), valley Munsu-tong, 20 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. This species is a single representative of the genus *Spilota* BURM. and a typical element of the Manchurian fauna.

***Proagopertha lucidula* (FALDERMANN, 1835)**

(Fig. 187)

Distribution. USSR — Ussuri, Primorskij Kraj (Chanka lake, Spassk), Manchuria (Mukden), North- and East China (MEDVEDEV, 1949; MACHATSCHKE, 1972; MEDVEDEV, 1974).

Material examined. 7 ♀♀. Prov. Phjǽngjang-si, Tesǽng-san (Mts, 124), 23 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), Phjǽngjang (86), 24 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), distr. Jongsǽng (41), Maram, 29 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Phjǽngan-namdo, distr. Sun-čhǽn (94), Džamo-ri, 27 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), distr. Sunan (104), Sǽkam-Čǽsudži (105), 21 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Hamgjǽng-pukto, distr. Kjǽngsǽng (44), Onpho-ri (77), 3 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. Beetles feed on leaves of apple-trees and other fruit-trees in orchards. They also occur in forest clearings on flowers, especially *Rosa davurica* PALL. Species new to the Korean fauna.

Mimela splendens (GYLLENHAL in SCHÖNHEER, 1817)

(Figs. 188, 189)

Distribution. USSR — Primorskij Kraj (Vladivostok), East China, Korea — Čedžu-do (Isl., 3), Japan — Honshu (Tochigi: Ootawara; Saitama-Tokorazawa, Hannǽ; Tokyo: Meguro, Kunitachi, Murayama; Kanagawa: Noborito; Niigata: Kurokawa, Imakawa, Shibata, Niigata; Shizuoka: Ohdaru spa; Aichi: Toyohashi; Gifu; Mie: Mt. Fujiwara, Iga, Yunoyama; Kyoto: Kyoto, Yamazaki; Osaka: Minoo; Hyogo: Hyonosen; Wakayama: Kanaya), Sado (Ogura-toge), Shikoku (Tokushima: Kamiryo; Ehime: Imabari, Mt. Ishizuchi), Kyushu (Fukuoka: Mt. Hikosan; Oita: Beppu; Kumamoto: Mt. Ichifusa; Kagoshima: Kurino spa, Kagoshima); Taiwan (Chiah sien, Fengshan, Suisharyo, Kangkou, Lienhwachi, Qixinsha, Puli, Nanshanchi), Indochina (Burma) (MEDVEDEV, 1949; NAKANE et BABA, 1960; MACHATSCHKE, 1972; NOMURA et KOBAYASHI, 1976).

Material examined. 2 ♂♂. Prov. Phjǽngan-namdo, distr. Kangsǽ (61), Thesǽng, vicinity of the salt lake, 18 VI 1974, Z. STEBNICKA (ISEZ); Prov. Hvanghe-namdo, Sujang-san (Mt., 121) vicinity of the water-fall, 3 VI 1974, Z. STEBNICKA (ISEZ).


Mimela chinensis KIRBY, 1823

(Figs. 190, 191)

Distribution. East China, Korea (MEDVEDEV, 1949; MACHATSCHKE, 1972).

Material examined. 1 ♂ and 1 ♀. Prov. Phjǽngjang-si, Jongak-san (Mt., 40), 29 V 1974, Z. STEBNICKA (ISEZ).

Remarks. Bionomics of the species of the genus *Mimela* KIRBY is little known; beetles fly in the daytime, feed on various plants. Both species were collected in the vicinity of waters in leafy brushwood.



Figs. 183—186. Male genitalia, lateral and dorsal view; 183, 184 — *Phyllopertha horticola* (L.); 185, 186 — *Spilota plagicollis* FAIRM.

Fig. 187. *Proagopertha lucidula* (FALD.) — last segment and claws of posterior tarsus. Figs. 188—194. Male genitalia, lateral and dorsal view; 188, 189 — *Mimela splendens* (GYLL.); 190, 191 — *M. chinensis* KIRBY; 192, 193 — *Rhombonyx holosericea* (FABR.); 194 — *Anomala* (*Chrysoplethisa*) *sieversi* HEYD.

Rhombonyx holosericea (FABRICIUS, 1787)

(Figs. 192, 193)

Distribution. European part of USSR to South Siberia, Amur, Primorskij Kraj, Sachalin; Mongolia, North China, Manchuria, Korea, Japan (MEDVEDEV, 1949; MACHATSCHKE, 1972; MEDVEDEV, 1974).

Material examined. 1 ♂ and 2 ♀♀. Prov. Hamgjǒng-pukto, distr. Kjǒng-sǒng (44), valley Onpho-ri (77), 11 IX 1966, H. SZEŁĘGIEWICZ and C. DZIA-DOSZ (ZIW); Prov. Janggang-do, distr. Počhǒn (90), vicinity of the village Posǒ-ri, 3 IX 1971, J. RAZOWSKI (ISEZ).

Remarks. Species linked to coniferous forests, expanded in taiga far to the west and crossed Ural. Beetles occur in woodland on sandy soil, in summer fly over forest clearings, sit on flowers and young trees, feed among others on pine needles. According to MEDVEDEV (1974) larvae destroy roots of garden plants. This is a frequent and numerous species in Primorskij Kraj. Duration of the developmental cycle not known yet.

Anomala (Chrysoplethisa) sieversi HEYDEN, 1887

(Fig. 194)

Distribution. North- and East China, Korea, Japan (MEDVEDEV, 1949; MACHATSCHKE, 1972).

Material examined. 61 ♂♂ and ♀♀. Prov. Kesǒng-si, vicinity of the water-fall Pakjǒn on the Čhǒnma-san (Mt., 13), 5—8 VI 1974, Z. STEBNICKA (ISEZ); Prov. Phjǒngan-namdo, distr. Sunan (104), Sǒkam-Čǒsudži (105), 21 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), distr. Kangsǒ (61), Thesǒng, 25—26 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), dist. Sunčhǒn (94), Džamo-ri, 27 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Phjǒngjang-si, distr. Jongsǒng (41), 29 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), distr. Samsǒk (106), Sǒngmun-ri, 22 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Kangvǒn-do, Čhǒnne (14), 10 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. Subgenus *Chrysoplethisa* REITT. is characteristic for the eastern part of Palaearctic. It forms a separate group of species ecology of which is little known. Fairly numerous specimens of the typical form *A. sieversi* and some specimens of ab. *atrocoerulea* REITT. were collected in Korea from flowers of bushes and herbaceous plants on sunny days in May and June. Similarly as other species of genus *Anomala* SAM., they probably feed on leaves and flowers of various plants, including cultivated ones.

Anomala (Euchronomala) cuprea (HOPE, 1839)

(Figs. 195, 196)

Distribution. USSR — Primorskij Kraj (Vladivostok), Kurile Islands (Ale-chino, Goriače Lake, Veslovskij Pen., Sernovodsk); North-Eastern China, Korea, Japan — Hokkaido Honshu (Kurokawa, Niigata, Hishizato), Sado (Matsugasaki, Mt. Donden), Shikoku, Kyushu (MEDVEDEV, 1949; NAKANE et BABA, 1960; MEDVEDEV et ERMOLENKO, 1969; MACHATSCHKE, 1972; KRYVOLUCKAJA, 1973; MEDVEDEV, 1974).

Material examined. 19 ♂♂ and ♀♀. Prov. Phjŏngan-namdo, distr. Sunčhŏn (94), Džamo-san (hills), 27 VIII 1971, J. PAWŁOWSKI (ISEZ), distr. Sunan (104), Sŏkam-Čŏsudži (105), 21 VIII 1971, A. SZEPTYCKI (ISEZ); Prov. Phjŏngjang-si, Jongak-san (Mt., 40), 20 VIII 1971, J. RAZOWSKI (ISEZ); Prov. Kangvŏn-do, Vŏnsan (118), 1 IX 1966, H. SZEŁĘGIEWICZ and C. DZIADOSZ (ZIW), 20 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), Kymgang-san (Mts, 53), 18 VI 1974, Z. STEBNICKA (ISEZ); Prov. Hamhyng-si, Tŏksan-ri 12 km N of Hamhyng (19), 14 IX 1966, H. SZEŁĘGIEWICZ and C. DZIADOSZ (ZIW).

Remarks. Species fairly numerous and widely distributed in Korea. Imagines feed on leaves and pollen of various plants, among others of roses (*Rosa rugosa* THUNB.). Larvae are enemies of corn and other cultivated plants.

***Anomala (Euchronomala) viridana* KOLBE, 1886**

Distribution. USSR — Primorskij Kraj (Vladivostok), Kurile Islands, (Kunašir, Alechino); Korea, Japan — Honshu (Kurokawa, Niigata, Araiama, Kinoto, Maki), Sado (Matsugasaki, Nakaoku), Shikoku (MEDVEDEV, 1949; NAKANE et BABA, 1960; MEDVEDEV et ERMOLENKO, 1969; MACHATSCHKE, 1972; KRYVOLUCKAJA, 1973).

Material examined. 5 ♀♀. Prov. Kangvŏn-do, Vŏnsan (118), 1 IX 1966, H. SZEŁĘGIEWICZ and C. DZIADOSZ (ZIW).

Remarks. Species very similar to the former one. Differs from it by slightly more mat surface of the upper side of body and more dense punctation of elytra.

***Anomala (Anomala) corpulenta* MOTSCHULSKY, 1853**

(Fig. 197)

Distribution. North China (Prov. Chebei-Tianxin), Korea, Japan (MEDVEDEV, 1949; MACHATSCHKE, 1972).


Material examined. 6 ♂♂ and ♀♀. Prov. Phjŏngjang-si, Phjŏngjang (86), 26 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Kangvŏn-do, Vŏnsan (118), 1 IX 1966, H. SZEŁĘGIEWICZ and C. DZIADOSZ (ZIW).

***Anomala (Anomala) luculenta* ERICHSON, 1847**

(Figs. 198, 199)

Distribution. USSR — Transbajkal Region, Amur, Primorskij Kraj; North China, Manchuria, Mongolia, Korea (MEDVEDEV, 1949; MACHATSCHKE, 1972; MEDVEDEV, 1974; 1976).

Material examined. 11 ♂♂ and ♀♀. Prov. Phjŏngan-namdo, distr. Sunan (104), Sŏkam-Čŏsudži (105), 2 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), distr. Kangsŏ (61), Thesŏng, 24 VIII 1971, A. SZEPTYCKI (ISEZ), distr.



Figs 195—206. Male genitalia, lateral and dorsal view; 195, 196 — *Anomala (Euchronomala) cuprea* (HOPE); 197 — *A. (Anomala) corpulenta* MOTSCH.; 198, 199 — *A. (A.) luculenta* ER.; 200, 201 — *Blitopertha (Exomala) pallidipennis* REITT.; 202 — *B. (E.) orientalis* (WATERH.); 203, 204 — *B. (Blitopertha) conspurcata* (HAR.); 205, 206 — *Adoretus (Lepadoretus) sinicus* BURM. Fig. 207. *A. (L.) tenuimaculatus* WATERH. — last segment and claws of female anterior tarsus

Sunčhön (94), Džamo-san (hills), 27 VIII 1971, J. PAWŁOWSKI (ISEZ); Prov. Kangvön-do, Vönsan (118), 1 IX 1966, H. SZEŁĘGIEWICZ and C. DZIADOSZ (ZIW).

Remarks. Manchurian element, linked to sandy and halfsandy soils; penetrates toward west, probably along river valleys. Imagines and larvae are enemies of orchards. Beetles come to light.

***Blitopertha (Exomala) pallidipennis* REITTER, 1903**

(Figs. 200, 201)

Distribution. USSR — Primorskij Kraj; Mongolia, Manchuria, North Korea (MEDVEDEV, 1949; MACHATSCHKE, 1972; MEDVEDEV, 1974; 1976).

Material examined. 1 ♂ and 2 ♀♀. Prov. Kangvön-do, Vönsan (118), 14 VI 1974, Z. STEBNICKA (ISEZ); Prov. Hamgjöng-namdo, distr. Hjangsan (27), Mjohjang-san (Mts), 3 VIII 1959, B. PISARSKI and J. PRÓSZYŃSKI (ZIW).

Remarks. Manchurian element of a narrow range. Beetles occur on various bushes and herbaceous plants, feed on flowers and young leaves. Larvae are enemies of young forest cultures.

***Blitopertha (Exomala) orientalis* (WATERHOUSE, 1875)**

(Fig. 202)

Distribution. Korea, Japan — Hokkaido (Hakodate), Honshu (Echigo: Kurokawa, Gokahama, Tanibama, Mt. Hishigatake, Ginzandaira, Yoshigahira), Sado (Suizu, Hatano, Mt. Donden, Aoneba-goe), Shikoku, Kyushu, Tsushima, Yakushima, Amami-Ōshima, Ogasawara, Niijima, Shikine, Kouzu, Hachijo, Torishima; Hawaiian Islands, North America (MEDVEDEV, 1949; NAKANE et BABA, 1960; NOMURA, 1966; 1969; MACHATSCHKE, 1972; MEDVEDEV, 1974).

Material examined. 6 ♂♂ and ♀♀. Prov. Hamgjöng-pukto, valley of river Poro-čhön 20 km NW of Kjöngsöng (44), 4 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Hamgjöng-namdo, distr. Hjangsan (27), Mjohjang-san (Mts), 3 VIII 1959, B. PISARSKI and J. PRÓSZYŃSKI (ZIW).

Remarks. The collected specimens are uniformly black with opalescent lustre. Representatives of this species are enemies of vineyards and plantations of sugar cane.

***Blitopertha (Blitopertha) conspurcata* (HAROLD, 1878)**

(Figs. 203, 204)

Distribution. USSR — South-Eastern Siberia (vicinity of Blagoveščensk), Primorskij Kraj (vicinity of Chanka Lake); North China, Korea, Japan — Honshu (Echigo: Kurokawa, Maramatsu-hama, Sasaguchi-hama, Kinoto) (MEDVEDEV, 1949; NAKANE et BABA, 1960; MACHATSCHKE, 1972).

Material examined. 134 ♂♂ and ♀♀. Prov. Kesöng-si, Čhönma-san (Mt., 13), 6 VI 1974, Z. STEBNICKA (ISEZ); Prov. Kangvön-do, Vönsan (118), 14 VI 1974, Z. STEBNICKA (ISEZ), Čhönne (14), 10 VI 1965, M. MROCZKOWSKI and A. RIE-

DEL (ZIW), Kymgang-san (Mts, 53), 16 VI 1974, Z. STEBNICKA (ISEZ); Prov. Hamhyng-si, Hyngnam (28), 16 VI 1965, 11 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), distr. Hamdžu (29), Hyngpong-ri, 12 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Hamgjông-namdo, distr. Hjangsan (27), Hjangam-ri, 16—21 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Phjôngjang-si, distr. Jongsông (41), Maram, 19 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. Numerous specimens of this species were collected from flowers of briar-rose; petalae were nearly completely eaten. Similarly as other species of the same genus, occurs in great masses.

Adoretini

Adoretus (Lepadoretus) sinicus BURMEISTER, 1855

(Figs. 205, 206)

Distribution. China, Korea, Japan — Honshu (Tsuruga), Kyushu (Nagasaki), Miyako, Ishigaki, Iriomote, Heteruma, Yonaguni; Taiwan, Indochina, Hawaiian Islands (MEDVEDEV, 1949; NOMURA, 1965, 1966; 1970; MACHATSCHKE, 1972; MEDVEDEV, 1974).

Material examined. 143 ♂♂ and ♀♀. Prov. Phjôngjang-si, distr. Jongsông (41), Maram, 29 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), distr. Sam-sôk (106), village Sôngmun-ri, 22 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Phjônggan-namdo, distr. Kangsô (61), Thesông, 19 IX 1966, H. SZEŁĘGIEWICZ and C. DZIADOSZ (ZIW), distr. Sunčhôn (94), Džamo-ri, 27 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Kangvôn-do, Vôsna (118), 1—2 IX 1966, H. SZEŁĘGIEWICZ and C. DZIADOSZ (ZIW), Ungčîn 25 km S of Vôsna (118), 18 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), Munčhôn (71) 12 km NW of Vôsna (118), 19 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW); Prov. Hvanghe-namdo, Sujang-san (Mt., 121) near Hedžu (18), 3 VI 1974, Z. STEBNICKA (ISEZ); Prov. Hamgjông-namdo, distr. Hjangsan (27), Hjangam-ri, 19—20 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), 25 IX 1966, H. SZEŁĘGIEWICZ and C. DZIADOSZ (ZIW); Prov. Hamhyng-si, Čojang (15) 15 km SW of Hamhyng (19), 25 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW); Prov. Kesông-si, Čhônma-san (Mt., 13), 8 VI 1974, Z. STEBNICKA (ISEZ).

Remarks. The representatives of the subgenus *Lepadoretus* REITT. mainly inhabit the Oriental Realm; in Palaearctic occur only two species. Their biology is little known. According to MEDVEDEV (1949) they are active in the evenings and nights and are enemies of fruit-trees, roses and grape-vines. *A. sinicus* is fairly frequent and numerous in D.P.R.K, especially in southern Provinces. In the evenings comes to light, however, it was collected mainly during the day on various *Rosaceae*.

Adoretus (Lepadoretus) tenuimaculatus WATERHOUSE, 1875

(Fig. 207)

Distribution. China, Korea Japan — Honshu (Echigo: Kurokawa, Sakasamaki, Mt. Kasuga), Sado (Kawaharada), Ryu-kyu Islands; Indochina (Burma, Vietnam), Java, Hawaiian Islands (MEDVEDEV, 1949; NAKANE et BABA, 1960; MACHATSCHKE, 1972; MEDVEDEV, 1974).

Material examined. 2 ♀♀. Prov. Phjôngjang-si, Jongak-san (Mt., 40), 21 VIII 1971, at light of UV lamps, J. RAZOWSKI (ISEZ).

Remarks. According to MEDVEDEV (1949, 1974) this species similarly to the former one is an enemy of fruit-trees, roses and grape-vines. No numerous populations were found in D.P.R.K.

*Cetoniinae**Goliathini**Rhomborrhina (Rhomborrhina) japonica* HOPE, 1841

(Figs. 209, 210)

Distribution. Central- and South China, Korea, Japan — Honshu (Shibata, Niigata, Mt. Yoneyama), Awa-shima, Sado (Sukunegi, Katabe, Kawamo, Mt. Donden, Ogura-toge, Suizu, Tassha, Matsugasaki), Shikoku, Kyushu, Tsushima, Ohshima, Niijima, Kouzu (NAKANE et BABA 1960; MEDVEDEV; 1964; NOMURA, 1969).

Material examined. 1 ♂ and 1 ♀. Prov. Kesöng-si, vicinity of the water-fall Pakjön on the Čhönma-san (Mt., 13), 5—8 VI 1974, Z. STEBNICKA (ISEZ).

Remarks. Nothing is known of the bionomy of representatives of the genus. Korean specimens were collected in young forest, among bush-like shrubs including *Aceraceae* (*Kalopanax pictum*).

Dicranocephalus adamsi PASCOE, 1863

(Fig. 208)

Distribution. Tibet („Maenia”), China (Sichuan), South Korea („Kosan”, „Hpouno-Soun-Ouen”), Indochina — Vietnam (Bao Lac) (MEDVEDEV, 1964).

Material examined. 1 ♀. Prov. Hvanghe-namdo, Sujang-san (Mt., 121), 3 VI 1974, Z. STEBNICKA (ISEZ).

Remarks. Genus *Dicranocephalus* BURM., discussed by POUILLAUDE (1914), MEDVEDEV (1964), KUROSAWA (1968) and MIKŠIĆ (1971) contains 8 species both Palaearctic and Oriental ones. Bionomy of these species is not known; they are probably linked to woodland and occur rarely. One female specimen has been collected by the author on the southern slope of a mountain in the young leafy forest containing *Ginkgo biloba*, *Quercus dentata*, *Q. aliena*, *Kalopanax pictum* and *Fagus* sp. The specimen is carbon black, without a sign of creme or white coating. POUILLAUDE (1914) already reported the occurrence of females coloured similarly.




Fig. 208. *Dieranocephalus adamsi* PASC. — female head. Figs. 209, 210. *Rhomborrhina* (*Rhomborrhina*) *japonica* HOPE; 209 — male genitalia, lateral view; 210 — clypeus. Figs. 211—216. Male genitalia, lateral and dorsal view; 211, 212 — *Cetonia* (*Eucetonia*) *magnifica* BALL.; 213, 214 — *Potosia* (*Liocola*) *brevitarsis* (LEWIS); 215, 216 — *P.* (*Colopotosia*) *aerata* ER.

Cetoniini

Cetonia (*Eucetonia*) *magnifica* BALLION, 1870

(Figs. 211, 212)

Distribution. South-Eastern Siberia (Amur — Blagoveščensk, Bureinskij Chrebet, Primorskij Kraj), North- and East China, Korea (MEDVEDEV, 1964).

Material examined. 42 ♂♂ and ♀♀. Prov. Hvanghe-namdo, Sujang-san (Mt., 121), 2 VI 1974, Z. STEBNICKA (ISEZ); Prov. Kesong-si, vicinity of the

water-fall Pakjön on the Čhönma-san (Mt., 13), 5 VI 1974, Z. STEBNICKA (ISEZ); Prov. Phjôngjang-si, Jongak-san (Mt., 40), 24 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), 29 V 1974, Z. STEBNICKA (ISEZ), distr. Jongsöng (41), Maram, 29 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), Tesöng-san (Mts, 124), 23 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Janggang-do, distr. Hjesan (26), vicinity of the village Döngha-ri, 2 IX 1971, J. PAWŁOWSKI (ISEZ), distr. Počhön (90), valley of river Karim-čhön 5 km S from Počhönbo, 1 IX 1971, J. PAWŁOWSKI (ISEZ); Prov. Kangvön-do, Kymgang-san (Mts, 53) near Kosöng (122), 16—18 VI 1974, Z. STEBNICKA (ISEZ); Prov. Hamgjöngnamdo, distr. Hjangsan (27), Hjangam-ri, 21 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), distr. Hongvön (30), Jönpong-ri, 8 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Čhöngdžin-si, mountain pass Musan-rjöng in the mountain chain Hamgjöng-sanmek, 2 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Hamhyng-si, distr. Hamdžu (29), Hyngpong-ri, 12 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Phjônggan-namdo, distr. Sunčhön (94), Džamo-ri, 27 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), distr. Sunan (104), Sökam-Čösudži (105), 21 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. Beetles occur in woodland, on large forest clearings and glades, in parks. They sit on flowers of various plants, in particular of *Spiraeoideae*, *Tamaricaceae* and *Caprifoliaceae*. Species widely spread in Korea, frequent but not numerous.

***Potosia (Liocola) brevitarsis* (LEWIS, 1879)**

(Figs. 213, 214)

Distribution. USSR — Chabarovskij and Primorskij Kraj, Sachalin, Kurile Islands (Vlc. Mendelev, Rejdovo, Kujbyševo); Mongolia, North-Eastern China, Taiwan, Korea, Japan — Hokkaido, Honshu (Kurokawa), Sado (Futami, Matuyama), Shikoku, Kyushu (NAKANE et BABA, 1960; MEDVEDEV, 1964; ENDRÖDI, 1971; KRYVOLUCKAJA, 1973; MEDVEDEV, 1974).

Material examined. 6 ♂♂ and ♀♀. Prov. Phjônggan-namdo, distr. Sunan (104), Sökam-Čösudži (105), 2 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), distr. Kangsö (61), Thesöng, 13 VI 1974, Z. STEBNICKA (ISEZ); Prov. Phjôngjang-si, Jongak-san (Mt., 40), 25 VIII 1966, H. SZEŁĘGIEWICZ and C. DZIADOSZ (ZIW), Tesöng-san (Mts, 124), 29 VIII 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), Mankjööngde, 31 VIII 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW); Prov. Hamhyng-si, distr. Hamdžu (29), Hyngpong-ri, 12 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. Beetles are enemies of fruit-trees, maize and other cultivated plants. No numerous population found in D.P.R.K.

Potosia (Calopotosia) aerata (ERICHSON, 1834)

(Figs. 215, 216)

Distribution. USSR — Primorskij Kraj (Vladivostok); East China, Korea, Japan — Honshu, Kyushu (MEDVEDEV, 1964).

Material examined. 2 ♂♂ and 1 ♀. Prov. Phjǽngjang-si, Jongak-san (Mt., 40), 29 V 1974, Z. STEBNICKA (ISEZ), distr. Jongsǽng (41), Maram, 29 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Potosia (Potosia) famelica (JANSON, 1879)

(Figs. 217, 218)

Distribution. USSR — Jevrejskaja A. O., Amur, Primorskij Kraj; East China, Korea (MEDVEDEV, 1964).

Material examined. 1 ♂ and 1 ♀. Prov. Phjǽngjang-si, Jongak-san (Mt., 40), 29 V 1974, Z. STEBNICKA (ISEZ); Prov. Kangvǽn-do, Samil-pho (123), vicinity of the lake, 19 VI 1974, J. PAWŁOWSKI (ISEZ).

Remarks. Species linked to leafy forests. According to MEDVEDEV (1964) and MIKŠIĆ (1959) in the southern part of the range occurs subspecies *P. famelica scheini* MIKŠ., which inhabits a part of Eastern China and South Korea including Čedžu-do (Isl., 3).

Poecilophilides rusticola (BURMEISTER, 1842)

(Figs. 219, 220)

Distribution. USSR — Amur, Ussuri, Daurija, Primorskij Kraj; Mongolia, Korea, North- and Central China, Japan — Honshu (Niigata), Kyushu, Sado (Kanazawa) (TESAŘ, 1959; NAKANE et BABA, 1960; MEDVEDEV, 1964).

Material examined. 95 ♂♂ and ♀♀. Prov. Phjǽngjang-si, Tesǽng-san (Mts, 124), 22 VIII 1966, H. SZEŁĘGIEWICZ and C. DZIADOSZ (ZIW), 29 VIII 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), 18 V 1974, Z. STEBNICKA (ISEZ), distr. Samsǽk (106), Sǽngmun-ri, 22 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).


Remarks. Species linked to leafy forests of a Manchurian type. Beetles fly from April to the beginning of September in lowlands and in mountains (up to 1000 m above sea level). They stay on flowers of various plants, are also found under bark of leafy trees. According to MEDVEDEV (1964) this species is frequent in Primorskij Kraj; very numerous locally in D.P.R.K.

Glycyphana (Glycyphana) fulvistemma (MOTSCHULSKY, 1860)

(Figs. 221, 222)

Distribution. USSR — East- and South Bajkal, Amur, Primorskij Kraj; Manchuria, North China, Korea, Japan — Honshu (Kurokawa); Taiwan (NAKANE et BABA, 1960; MEDVEDEV, 1964; MIKŠIĆ, 1970; MEDVEDEV, 1974).

Material examined. 4 ♂♂ and ♀♀. Prov. Phjôngan-namdo, distr. Sunčhôn (94), Džamo-ri, 27 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Phjôngjang-si, distr. Jongsông (41), 29 V 1965, M. MROCZKOWSKI and A. RIE-



Figs. 217—226. Male genitalia, lateral and dorsal view; 217, 218 — *Potosia (Potosia) famelica* (JANS.); 219, 220 — *Poecilophilides rusticola* (BURM.); 221, 222 — *Glycyphana* (*Glycyphana*) *fulvistemma* (MOTSCH.); 223, 224 — *Oxycetonia jucunda jucunda* (FALD.); 225, 226 — *Clinterocera mandarina* (WESTW.)

DEL (ZIW); Prov. Hamgjông-namdo, distr. Hjangsan (27) valley Manphok-tong in the Mjohjang-san (Mts), 29 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. Manchurian species, occurs in woodland. Beetles fly from Spring to Autumn; feed on flowers of various plants, among others on fruit-trees. Numerous populations were not found in D.P.R.K.

Oxycetonia jucunda jucunda (FALDERMANN, 1835)

(Figs. 223, 224)

Distribution. North-and East India, Tibet, Nepal, North-and Central China, Manchuria, USSR — Kamchatka Pen., Sachalin, Kurile Islands, Komandor Islands, Amur, Ussuri, Primorskij Kraj; Korea — Čedžu-do (Isl., 3); Japan — Hokkaido, Honshu (Kurokawa, Shibata, Niigata, Kakuda-hama, Noo, Miomote, Yoshigahira), Awa-shima, Sado (Futami, Ogura-toge, Ookura-goe), Shikoku, Kyushu, Tsushima, Yakushima, Tanegashima, Ohshima, Toshima, Niijima, Shikine, Kouzu, Miyake, Hachijo, Aogashima; Central and North America (introduced) (NAKANE et BABA, 1960; MEDVEDEV, 1964; NOMURA, 1969; MEDVEDEV, 1974).

Material examined. 546 ♂♂ and ♀♀. Prov. Kangvön-do, Kymgang-san Mts, 53), 16—18 VI 1974, Z. STEBNICKA (ISEZ), Samil-pho (123), vicinity of the lake, 19 VI 1974, J. PAWŁOWSKI and A. SZEPTYCKI (ISEZ), Vönsan (118), 20 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), 2 IX 1966, H. SZEŁĘGIEWICZ and C. DZIADOSZ (ZIW), Sičung-ho (107), 18 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), Munčhön (71), 12 km NW of Vönsan (118), 19 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW); Prov. Kesöng-si, vicinity of the water-fall Pakjön on the Čhönma-san (Mt., 13), 6—8 VI 1974, Z. STEBNICKA (ISEZ); Prov. Phjöngjang-si, Jongak-san (Mts, 40), 29 V 1974, Z. STEBNICKA (ISEZ), 25 VIII 1966, H. SZEŁĘGIEWICZ and C. DZIADOSZ (ZIW), Tesöng-san (Mts, 124), 23 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), 28 VIII 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), Ponghva-ri (88), 24 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), distr. Jongsöng (41), Maram, 29 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), Mankjööngde, 31 VIII 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW); Prov. Hvanghe-namdo, Sujang-san (Mt., 121), 2 VI 1974, Z. STEBNICKA (ISEZ); Prov. Phjöng-gan-namdo, distr. Sunčhön (94), Džamo-ri, 27 VIII 1971, J. PAWŁOWSKI (ISEZ), 27 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), distr. Kangsö (61), The-söng, 23 VIII 1971, A. SZEPTYCKI (ISEZ), 19 IX 1966, H. SZEŁĘGIEWICZ and C. DZIADOSZ (ZIW), 26 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), distr. Sunan (104), Sökam-Čösudži (105), 21 V 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), 2 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), Junha-ri, 13 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), Mjohjang-san (Mts), 24 IX 1966, H. SZEŁĘGIEWICZ and C. DZIADOSZ (ZIW); Prov. Janggang-do, distr. Hjesan (26), vicinity of the village Dönggha-ri, 2 IX 1971, J. RAZOWSKI (ISEZ); Prov. Hamgjöng-pukto, distr. Kjööngsöng (44), valley Onpho-ri (77), 23—24 V 1974, J. PAWŁOWSKI (ISEZ), 3 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), 6 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW); Prov. Hamhyng-si, Čojang (15), 25 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), Oro (78), 15 IX 1966, H. SZEŁĘGIEWICZ and C. DZIADOSZ (ZIW), Tök-san-ri, 14 IX 1966, H. SZEŁĘGIEWICZ and C. DZIADOSZ (ZIW), distr. Hamdžu (29), Hyngpong-ri, 12 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov.

Hamgjöng-namdo, distr. Hongvön (30), Jönpong-ri, 8 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), distr. Hjangsan (27), Hjangam-ri, 21 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), Sangvön-am (109), 17 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Čhöngdžin-si, distr. Purjöng (87), Thomak-tong, 8 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW).

Remarks. Species is characterised by a great individual variability. Occur in a number of variants. Moreover, it forms geographic races reported from Japan by NOMURA (1966). According to their range (northern or southern) beetles fly in Spring or Summer in open woodland, in lowlands or mountains. They feed on flowers of various trees, bushes and herbaceous plants, are enemies of orchards in blossom. The species occurs in great masses in Primorskij Kraj, however, its number decreases gradually to the direction of north. In North Korea is widely distributed and numerous. Occurrence in great masses (10—20 specimens on a single inflorescence) was found locally. Besides of a typical form, variants *ab. ferruginosa* REITT., *ab. vitticollis* REITT., *ab. marginalis* MEDV., *ab. kuperi* SCHAUM, *ab. dolens* KR. occur frequently.

Cremastochilini

Clinterocera mandarina (WESTWOOD, 1874)

(Figs. 225, 226)

Distribution. USSR — Primorskij Kraj (Spassk, Jakovlevka, Duškino, Sučan), North China, Korea (MEDVEDEV, 1964).

Material examined. 2 ♂♂. Prov. Hamgjöng-pukto, distr. Kjöngsöng (44), Jonghen ad Džuyr (17), 25 V 1974, Z. STEBNICKA (ISEZ); Prov. Hvanghe-namdo, vicinity of the water-fall Sujang-san (Mt., 121) near Hedžu (18), 3 VI 1974, Z. STEBNICKA (ISEZ).

Remarks. The way of life of the representatives of the tribe *Cremastochilini* does not differ from this of other *Cetoniinae*, however, their ecology is little known. Beetles are day-active, occur in Spring and Summer. *C. mandarina* is one of two species that inhabit Korean Peninsula; found sporadically in the leafy forests of a Manchurian type.

Valginae


Dasyvalgus angusticollis (WATERHOUSE, 1875)

(Figs. 227, 228)

Distribution. Korea, Japan — Hokkaido, Honshu (Echigo: Kurokawa, Kuramitsu, Kamiishikawa, Meiji-mura, Miomote), Awa-shima, Sado (Matsugasaki, Nakaoku, Mt. Kinpoku), Shikoku, Kyushu, Tsushima, Yakushima, Tanegashima, Ohshima, Toshima, Nijima, Miyake (NAKANE et BABA, 1960; MEDVEDEV, 1964; NOMURA, 1969).

Material examined. 6 ♂♂ and ♀♀. Prov. Kesöng-si, vicinity of the waterfall Pakjön on the Čhönma-san (Mt., 13) 5—8 VI 1974, Z. STEBNICKA (ISEZ).

Remarks. Beetles dendrophilic, fly during warm sunny days, occur on leaves and flowers of trees and bushes.



Figs. 227—232. Male genitalia, lateral and dorsal view; 227, 228 — *Dasyvalgus angusticollis* (WATERH.); 229, 230 — *Lasiotrichius succinctus* (PALL.); 231, 232 — *Gnorimus subopacus* MOTSCH.

Trichiinae

Trichiini

Lasiotrichius succinctus (PALLAS, 1781)

(Figs. 229, 230)

Distribution. USSR — Central- and East Siberia, Bajkal, Sachalin, Southern Kurile, Primorskij Kraj, North Mongolia, Manchuria, North- and East China, Korea, Japan — Honshu (Echigo: Renge-spa), Hokkaido, Shikoku, Kyushu, Ryu-kyu Islands (NAKANE et BABA, 1960; MEDVEDEV, 1960; TESAR 1968; ENDRÖDI, 1971; KRYVOLUCKAJA, 1973).

Material examined. 80 ♂♂ and ♀♀. Prov. Kangvön-do, Kymgang-san (Mts, 53), 18 VI 1974, Z. STEBNICKA (ISEZ), Samil-pho (123) vicinity of the lake, 19 VI 1974, J. PAWŁOWSKI (ISEZ), Masin-rjöng, mountain pass, 34 km W of

Võnsan (118), 16 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW); Prov. Hvanghe-namdo, vicinity of the water-fall Sujang-san (Mt., 121) near Hedžu (18), 3 VI 1974, Z. STEBNICKA (ISEZ); Prov. Janggang-do, distr. Hjesan (26), Dõngha-ri, 2 IX 1971, J. PAWŁOWSKI (ISEZ); Prov. Phjõngjang-si, Jongak-san (Mts, 40) 29 V 1974, Z. STEBNICKA (ISEZ); Prov. Kesõng-si, vicinity of the water-fall Pakjõn on the Čhõnma-san (Mt., 13), 7 VI 1974, Z. STEBNICKA (ISEZ); Prov. Hamgjõng-pukto, distr. Kjõngsõng (44), valley Onpho-ri (77), 11 IX 1966, H. SZELEGIWICZ and C. DZIADOSZ (ZIW), 5 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW), Džuyt (17), 7 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW); Prov. Hamgjõng-namdo, distr. Hjangsan (27), Hjangam-ri, 22 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW), Sangvõn-am (109), 17 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW); Prov. Hamhyng-si, distr. Hamdžu (29), Hyngpong-ri, 12 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. East-Siberian species, occurs in lowlands and mountains, in leafy and coniferous forests. Beetles are active from Spring to early Autumn, feed on flowers of herbaceous plants, trees and bushes. Species frequent and numerous in D.P.R.K., found mainly on large inflorescences of *Rosaceae* and *Caprifoliaceae*.

Gnorimus subopacus MOTSCHULSKY, 1860

(Figs. 231, 232)

Distribution. USSR — Amur, Ussuri, Primorskij Kraj, Sachalin, Kurile Islands (Kunašir); North-Eastern China, Manchuria, Korea, Japan (MEDVEDEV, 1960; MEDVEDEV et ERMOLENKO, 1969; KRYVOLUCKAJA, 1973).

Material examined. 2 ♂♂. Prov. Kangvõn-do, Kymgang-san (Mts, 53), 16 VI 1974, Z. STEBNICKA (ISEZ); Prov. Hamgjõng-pukto, Nadžin (74), 5 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

Remarks. Manchurian species, linked to leafy and mixed forests. Beetles are active in Summer, feed on flowers of bushes and herbaceous plants. The Korean specimens have been collected on the flowers of *Magnoliaceae*.

Trogidae

Trox sabulosus (LINNAEUS, 1758)

(Fig. 102)

Distribution. Europe, Caucasus, Siberia, Korea — Čedžu-do (Isl., 3) (HORION, 1958 et all.).

Material examined. 1 ♂. Prov. Hamgjõng-namdo, distr. Hjangsan (27), Hjangam-ri, 18 VI 1965, M. MROCZKOWSKI and A. RIEDEL (ZIW).

*Lucanidae**Psalidoremus inclinatus* (MOTSCHULSKY, 1857)

Distribution. Taiwan, Korea, Japan — Hokkaido, Honshu, Sado, Shikoku, Kyushu, Yakushima, Kuchinoerabu, Ohshima, Nijima, Kouzu, Miyake, Mikura (DIDIER et SÉGUY, 1952—1953; BENESH, 1960; NOMURA, 1965; 1966; NISHIO, 1970; NAKANE, 1977 — P. 27, Fig. 577).

Material examined. 1 ♀. Prov. Hamgjong-pukto, distr. Kjongssong (44), vicinity of Dzuyr (17), 7 IX 1970, R. BIELAWSKI and M. MROCZKOWSKI (ZIW).

Macrodorcas rubrofemoratus (VOLLENHOVEN, 1865)

Distribution. USSR — Primorskij Kraj, Kurile Islands (Alechino); North- and Central China, Taiwan, Korea, Japan — Hokkaido, Honshu, Shikoku, Kyushu, Amami-Oshima (DIDIER et SÉGUY, 1952—1953; BENESH, 1960; NOMURA, 1966; NISHIO, 1970; KRYVOLUCKAJA, 1973; NAKANE, 1977 — Pl. 27, Fig. 579).

Material examined. 1 ♀. Prov. Kangvön-do, Kymgang-san (Mts, 53) near Kosong (122), 18 VI 1974, at light, Z. STEBNICKA and A. SZEPTYCKI (ISEZ).

Macrodorcas striatipennis MOTSCHULSKY, 1861

Distribution. USSR — Kurile Islands (Alechino, Malokurilsk), Manchuria, North China, Taiwan, Korea, Japan — Hokkaido, Honshu, Sado, Shikoku, Kyushu, Yakushima, Tanegashima, Tsushima (DIDIER et SÉGUY, 1952—1953; BENESH, 1960; NOMURA, 1969; KRYVOLUCKAJA, 1973; NAKANE, 1977 — Pl. 27, Figs. 584—586).

Material examined. 1 ♂. Prov. Kesong-si, vicinity of the water-fall Pakjon on the Chonma-san (Mt., 13), 8 VI 1974, Z. STEBNICKA (ISEZ).

Serrognathus titanus (BOISDUVAL, 1835)

Distribution. Korea, Japan — Honshu, Shikoku, Kyushu, Yakushima, Nakanoshima, Ohshima, Toshima, Nijima, Iki; Taiwan, Central- and South China, East India, Assam, Indochina, Philippines, Sunda Islands (DIDIER et SÉGUY, 1952—1953; BENESH, 1960; NOMURA, 1965; 1966; 1969; NAKANE, 1977 — Pl. 27, Figs. 587, 588).

Material examined. 4 ♂♂. Prov. Hvangde-namdo, Hedzu (18), 1 VI 1974, at light, Z. STEBNICKA (ISEZ); Prov. Kangvön-do, Kymgang-san (Mts, 53) near Kosong (122), 18 VI 1974, at light, Z. STEBNICKA (ISEZ).

Dorcus hopei (SAUNDERS, 1854)

Distribution. Korea, Manchuria, North-, Central- and South China, Taiwan, Japan, Indochina (DIDIER et SÉGUY, 1952—1953; BENESH, 1960; NAKANE, 1977 — Pl. 28, Figs. 600, 601).

Material examined. 1 ♀. Prov. Kangvŏn-do, Kymgang-san (Mts, 53) near Kosŏng (122), 19 VI 1974, at light Z. STEBNICKA and A. SZEPTYCKI (ISEZ).

Dorcus montivagus (LEWIS, 1883)

Distribution. North China, Korea, Japan — Hokkaido, Honshu, Shikoku, Kyushu, Ohshima (DIDIER et SÉGUY, 1952—1953; BENESH, 1960; NOMURA 1969; NAKANE, 1977 — Pl. 28, Figs. 602, 603).

Material examined. 1 ♀. Prov. Kangvŏn-do, Kymgang-san (Mts, 53) near Kosŏng (122), 18 VI 1974, at light, Z. STEBNICKA and A. SZEPTYCKI (ISEZ).

Institute of Systematic and Experimental Zoology
Polish Academy of Sciences,
Ślawkowska 17, 31-016 Kraków, Poland

REFERENCES

- ALECHIN V. V. 1951. Rastitelnost SSSR v osnovnykh zonakh. Moskva—Leningrad, 511 pp.
- ALLIBERT A. 1847. Note sur divers Insectes Coléoptères trouvés dans des graines rapportées de Couton. Rev. Zool., **10**: 11—19.
- ARROW G. J. 1913. Notes on the Lamellicorn *Coleoptera* of Japan and descriptions of a few new species. Ann. Mag. nat. Hist., **8** (12): 394—408.
- BALLION E. 1871. Eine centurie neuer Käfer aus der Fauna des russischen Reiches. Bull. Soc. imp. Nat. Mosc., **43** (2): 320—353.
- BALTHASAR V. 1932. *Aphodius haroldianus* n. n. für *Aphodius apicalis* Har. Ent. NachrBl., **6**: 1—7.
- BALTHASAR V. 1933. Neue *Aphodius*-Arten aus dem palaearktischen Asien. Koleopt. Rdsch., **19**: 139—146.
- BALTHASAR V. 1935. Einige neue palaearktische Scarabaeiden. Ent. Blätt., **31** (3): 120—124.
- BALTHASAR V. 1938. Neue palaearktische *Aphodius*-Arten. Čas. čsl. Spol. ent., **35**: 6—13.
- BALTHASAR V. 1963—1964. Monographie der *Scarabaeidae* und *Aphodiidae* der palaearktischen und orientalischen Region. *Coleoptera: Lamellicornia*, **1**, 391 pp.; **2**, 627 pp.; **3**, 652 pp.
- BALTHASAR V. 1965. Neue *Aphodius*-Arten aus Nepal. 122. Beitrag zur Kenntnis der *Scarabaeoidea*. Khumbu Himal, **2**: 108—113.
- BATES H. W. 1888. On a collection of *Coleoptera* from Korea (Tribes: *Geodephaga*, *Lamellicornia* and *Longicornia*) made by Mr. J. H. LEECH, F. Z. S. Proc. zool. Soc. Lond., 367—380.
- BENESH B. 1960. *Lucanidae*. In: *Coleopterorum Catalogus* (JUNK-SCHENKLING), Suppl., **8**, 178 pp.

- BOISDUVAL J. B. 1835. Faune entomologique de l'Océan Pacifique. Voyage l'Astrolabe, Coléopt., 2, 6: 267—716.
- BOUCOMONT A. 1924. Synonymies and homonymies de Lamellicornes Coprophages (Col.). Bull. Soc. ent. Fr., 114—115.
- BRENSKE E. 1892. Neue Arten der Coleopteren Gattung *Holotrichia* (*Lachnosterna*). Berl. ent. Z., 37: 159—192.
- BRENSKE E. 1897. Die *Serica*-Arten der Erde, I. Berl. ent. Z., 42: 345—438.
- BRULLÉ A. 1832. Expédition scientifique de Morée. Section des Sciences Physiques. Tome III., 1-re Partie. Zoologie. Deuxième Section. — Des animaux articulés. Paris, 404 pp.
- BURMEISTER H. 1842. Handbuch der Entomologie, III. Berlin, 135—638, 769—812.
- BURMEISTER H. 1855. Handbuch der Entomologie. 2. *Lamellicornia Phyllophaga Chaenochela*. Berlin, 10+570 pp.
- CHANG YOU-WEI. 1965. Revision of the Chinese may beetles of the genus *Holotrichia* HOPE (*Coleoptera*, *Scarabaeidae*). Acta zootax. sin., 2 (1): 37—56.
- CZEKANOWSKI J. 1930. Zarys antropologii Polski. Lwów, 592 pp.
- DIDIER R., SÉGUY E. 1952—1955. Catalogue illustré des Lucanides du Globe. Encyclopédie entomologique, Paris, 27, 223 pp. 28, 112 tbl.
- ENDRÖDI S. 1965. 62. *Lamellicornia* der II Expedition. Ergebnisse der zoologischen Forschungen von Dr. Z. KASZAB in der Mongolei (*Coleoptera*). Reichenbachia, 7 (21): 191—199.
- ENDRÖDI S. 1967. 103. *Lamellicornia* der III Expedition. Ergebnisse der zoologischen Forschungen von Dr. Z. KASZAB in der Mongolei (*Coleoptera*). Reichenbachia, 9 (15): 135—144.
- ENDRÖDI S. 1968. 138. *Lamellicornia* der IV Expedition. Ergebnisse der zoologischen Forschungen von Dr. Z. KASZAB in der Mongolei (*Coleoptera*). Reichenbachia, 11 (8): 69—81.
- ENDRÖDI S. 1969. 174. *Lamellicornia* der V Expedition. Ergebnisse der zoologischen Forschungen von Dr. Z. KASZAB in der Mongolei (*Coleoptera*). Reichenbachia, 11 (28): 287—299.
- ENDRÖDI S. 1971. 201. *Lamellicornia* der VI Expedition. Ergebnisse der zoologischen Forschungen von Dr. Z. KASZAB in der Mongolei (*Coleoptera*). Faun. Abh., 3 (11): 111—129.
- ERICHSON W. F. 1834. MEYENS Beiträge zur Zoologie gesammelt auf einer Reise um die Erde, u. W. ERICHSONS u. H. BURMEISTERS Beschreibungen u. Abbildungen der auf dieser Reise gesammelten Insecten. Acta Acad. Leopold. Carol., 16 (1): 219—284.
- ERICHSON W. F. 1847. Naturgeschichte der Insecten Deutschlands. *Coleoptera*, 3, 1, 3. Berlin, 481—800.
- ESCHSCHOLTZ J. F. 1818. Decades tres Eleutheratorum novorum. Mém. Acad. Sci. St. Pétersb., 6: 451—484.
- FABRICIUS J. Ch. 1775. Systema Entomologiae, sistens Insectorum classes, ordines, genera, species, adiectis synonymis, locis, descriptionibus, observationibus. Flensburgi et Lipsiae, 32+832 pp.
- FABRICIUS J. Ch. 1787. Mantissa Insectorum. Hafniae, 1, 382 pp.
- FABRICIUS J. Ch. 1798. Supplementum entomologiae systematicae. Hafniae, 572 pp.
- FAIRMAIRE L. 1886. Descriptions de Coléoptères de l'intérieur de la Chine. Ann. Soc. ent. Fr., 6 (6): 303—356.
- FALDERMANN F. 1835. Coleopterorum ab ill. BUNGIO in China boreali, Mongolia et montibus Altaicis collectorum, nec non ab ill. TURCZANINOFFIO et STSCHUKINO e provincia Irkutsk missorum illustrationes. Mém. Acad. Sci. St. Pétersb., 2: 337—464.
- GEBLER F. 1832. Notice sur les Coléoptères qui se trouvent dans le district des mines de Nertschinsk, dans la Sibérie orientale. Nouv. Mém. Soc. imp. Mosc., 2: 23—78.
- GOOD R. 1974. The geography of the flowering plants. London, 557 pp.
- HAROLD E. v. 1861. Beiträge zur Kenntnis einiger coprophagen Lamellicornien. II Stück. Berl. ent. Z., 5: 92—115.
- HAROLD E. V. 1862. Beiträge zur Kenntnis einiger coprophagen Lamellicornien. III Stück. Berl. ent. Z., 6: 138—171.
- HAROLD E. v. 1870. Ueber Nomenclatur. I Stück. Col. Hefte, 6: 37—69.

- HAROLD E. v. 1874. Beiträge zur Kenntnis einiger coprophagen Lamellicornien. VIII Stück. Berl. ent. Z., **18**: 177—208.
- HAROLD E. v. 1878. Beiträge zur Käferfauna von Japan. Dt. ent. Z., **22**: 65—88.
- HAROLD E. v. 1879. Neue Amur-Käfer. Dt. ent. Z. **23**: 145—146.
- HAROLD E. v. 1886. Coprophage Lamellicornien beschrieben. Berl. ent. Z., **30**: 141—149.
- HERBST J. F. W. 1789. Natursystem aller bekannten in- und ausländischen Insecten... Käfer, **2**, 330 pp.
- HEYDEN L. v. 1886. Die Coleopteren-Fauna des Suyfun-Flusses. Dt. ent. Z., **30**: 269—277.
- HEYDEN L. v. 1887. Vierter Beitrag zur Kenntnis der Coleopteren-Fauna der Amurländer. Dt. ent. Z., **31**: 297—304.
- HOPE F. W. 1839. Monograph on the Coleopterous genus *Euchlora* MC LEAY. Proc. zool. Soc. Lond., **7**: 65—75.
- HOPE F. W. 1841. Description of some new Lamellicorn *Coleoptera* from Northern India. Trans. R. ent. Soc. Lond., **3**: 62—67.
- HORION A. 1958. Faunistik der mitteleuropäischen Käfer. Band VI: *Lamellicornia* (*Scarabaeidae* — *Lucanidae*). Überlingen-Bodensee, XXIII + 343 pp.
- JACOBSON G. 1914. De Hopliinis novis palaearticeis (*Coleoptera*, *Scarabaeidae*). Russk. ent. Obozr., **14**: 1—5.
- JANSON O. E. 1879. Notices of new or little known *Cetoniidae*. Cist. Ent., **2**: 537—539.
- KALININA O. I. 1977. Revizija chrušej roda *Holotrichia* HOPE (*Col. Scarab.*) fauny SSSR. Ent. Obozr., **56** (4): 788—794.
- KIRBY W. F. 1823. A description of some Insects which appear to exemplify Mr W. S. MAC LEAYS doctrine of affinity and analogy. Trans. R. ent. Soc. Lond., **14** (1): 93—110.
- KOLBE H. J. 1886. Beiträge zur Kenntnis der Coleopteren-Fauna Koreas, bearbeitet auf Grund der von Herrn Dr. C. GOTTSCHKE während der Jahre 1883 u. 1884 in Korea veranstalteten Sammlung... Arch. Naturgesch., **52** (1): 139—240.
- KOMAROV V. L. 1949. Flora Mančžurii. Izbr. soč., **2**, 55 pp.
- KÔNO H. 1937. Die Lamellicornien aus den Kurillen. II. Insecta matsum. **12**: 6—8.
- KONTKANEN P. 1956. Zur Kenntnis der palaearktischen Käferfauna. I. Ann. ent. fenn., **22** (2): 55—63.
- KOSTROWICKI A. S. 1965. Regionalizacja zoogeograficzna Palearktyki w oparciu o faunę motyli tzw. większych (*Macrolepidoptera*). Pr. geogr., **41**, 99 pp.
- KOSTROWICKI A. S. 1969. Geography of the Palearctic *Papilionidea* (*Lepidoptera*). PWN, Kraków, 380 pp.
- KRAATZ G. 1892. Monographische Revision der Ruteliden-Gattung *Popillia* SERV. Dt. ent. Z., 225—306.
- KRIŠTOFOVIČ A. N. 1932. Geologičeskij obzor strany Dalnego Vostoka. Leningrad—Moskwa, 290—297.
- KRIŠTOFOVIČ A. N. 1958. Proischoždenie flory Angarskoj suši. Mater. po ist. flory i rastitelnosti SSSR, **3**: 7—41.
- KRYVOLUCKAJA G. O. 1973. Entomofauna Kurilskich ostrovov. Osnovnye čerti i proischoždenie. Leningrad, 313 pp.
- KURENCOV A. I. 1952. Čto takoe mančžurskaja fauna? Trudy Leningr. obšč. estestvoispit., **21** (4): 214—237.
- KURENCOV A. I. 1960. Proischoždenie i osnovnye etapy razvitija mančžurskoj entomofauny. Tez. dokl. 4-go sezda Vsesojuzn. entomol. obšč., 73—76.
- KURENCOV A. I. 1961. Životnyj mir Dalnego Vostoka. Sb. „Dalnyj Vostok“, Moskwa—Leningrad, 246—285.
- KURENCOV A. I. 1974. Zoogeografija Dalnego Vostoka SSSR na primere rasprostranjenja češuekrylych — *Rhopalocera*. Novosibirsk, 156 pp.
- KUROSAWA Y. 1968. Notes on the Formosan Cetoniid Beetles. I — A. Revision on the For-

- mosan Species of the Genus *Dicranocephalus* WESTWOOD. Bull. nat. Sci. Mus., **11** (3): 225—234.
- LEWIS G. 1879. On certain New Species of *Coleoptera* from Japan. Ann. Mag. nat. Hist., **5** (4): 459—467.
- LEWIS G. 1883. On the *Lucanidae* of Japan. Trans. R. ent. Soc. Lond., **14**: 333—342.
- LEWIS G. 1895. On the Lamellicorn *Coleoptera* of Japan. Ann. Mag. nat. Hist., **6** (16): 374—408.
- LINNAEUS C. 1758. Systema Naturae per regna tria naturae, secundum Classes, Ordines, Genera, Species, cum characteribus, differentiis, synonymis, locis. Editio Decima, Reformata. Tomus I. Holmiae, IV + 824 pp.
- MACHATSCHKE J. W. 1972. *Scarabaeoidea: Melolonthidae, Rutelinae*. In: *Coleopterorum Catalogus* (JUNK-SCHENKLING), Suppl., **66**, 1, 361 pp.
- MANNERHEIM C. G. 1849. Insectes Coléoptères de la Sibérie orientale nouveaux ou peu connus. 2. Bull. Soc. imp. Nat. Mosc., **22** (1): 220—249.
- MANNERHEIM C. G. 1852. Zweiter Nachtrag zur Käferfauna der Nord-Amerikanischen Länder des Russischen Reiches. Bull. Soc. imp. Nat. Mosc., **25** (4): 283—387.
- MATSUMURA S. 1936. New *Caccobius*-Species in Japan with a tabular key. Insecta matsum., **11**: 61—66.
- MATSUMURA S. 1937. The Onthophagid Insects from Korea with Description of New Species. Insecta matsum., **12**: 1—6.
- MEDVEDEV S. I. 1949. Platinčatousye (*Scarabaeidae*), podsem. *Rutelinae* (chlebnye žuki i blizkie gruppy). Fauna SSSR. Žestkokrylye. X. Moskva—Leningrad, **3**, 371 pp.
- MEDVEDEV S. I. 1951—1952. Platinčatousye (*Scarabaeidae*), podsem. *Melolonthinae*, I—II. (chrušči). Fauna SSSR. Žestkokrylye. X. Moskva—Leningrad, **1**, 512 pp., **2**, 274 pp.
- MEDVEDEV S. I. 1960. Platinčatousye (*Scarabaeidae*), podsem. *Euchirinae, Dynastinae, Glaphyrinae, Trichiinae*. Fauna SSSR. Žestkokrylye. X. Moskva—Leningrad, **4**, 397 pp.
- MEDVEDEV S. I. 1964. Platinčatousye (*Scarabaeidae*), podsem. *Cetoniinae, Valginae*. Fauna SSSR. Žestkokrylye. X. Moskva—Leningrad, **5**, 374 pp.
- MEDVEDEV S. I. 1974. 7. Sem. *Scarabaeidae* — platinčatousye. In: Nasekomye i klešči — vrediteli sel'skochozjaistvennykh kultur. Žestkokrylye. Leningrad, **2**: 18—60.
- MEDVEDEV S. I. 1976. Platinčatousye (*Coleoptera, Scarabaeidae*), sobrannye entomologičeskim otrjadom Sovetsko-Mongolskoj biologičeskoj ekspedicii v 1970—1971 gg. In: Nasekomye Mongolii, **4**: 155—164.
- MEDVEDEV S. I., ERMOLENKO V. M. 1969. Materialy k faune nadsemejstva platinčatousykh žukov (*Coleoptera, Lamellicornia*) Kuril'skikh ostrovov (po materialam pervoj kuril'skoj ekspedicii AN USSR v 1965 g.). Zbirn. Prac zool. Muz., **33**: 61—64.
- MIKŠIĆ R. 1959. Beitrag zur Kenntnis der palaearktischen Arten der Gattung *Potosia* MULS. III. Acta Mus. maced. Sci. nat., **6**: 27—51.
- MIKŠIĆ R. 1970. Katalog der Gattung *Glycyphana* BURM. (*Col.*) 60. Beitrag zur Kenntnis der Scarabaeiden. Folia ent. hung., **23** (6): 141—147.
- MIKŠIĆ R. 1971. Einige Bemerkungen über die systematische Stellung und Einteilung der asiatischen *Goliathina* (78. Beitrag zur Kenntnis der *Scarabaeidae*). Acta ent. Jugosl., **7** (2): 37—47.
- MIWA Y. 1930. An enumeration of the Coprophagid-*Coleoptera* from Formosa with a table of the geographical distribution. Insecta matsum., **4**: 163—180.
- MOSER J. 1915. Neue *Serica*-Arten. Berl. ent. Z., **57**: 337—393.
- MOTSCHULSKY V. 1853—1854. Coléoptères du Nord de la Chine. Études Entomologiques, Helsingfors, **2**, 56 pp., **3**, 64 pp.
- MOTSCHULSKY V. 1857. Études Entomologiques, Helsingfors, **6**, 112 pp.
- MOTSCHULSKY V. 1860. Études Entomologiques, Helsingfors, **9**, 42 pp.
- MOTSCHULSKY V. 1860. Coléoptères rapportés de la Sibérie orientale et notamment des pays situés sur les bords du fleuve Amour par M. M. SCHRENCK, MAACK, DITMAR, VOZNESSEN-

- SKI etc. déterminés et décrits. In: Dr. L. v. SCHRENCK's Reisen und Forschungen im Amur-Lande. *Coleoptera*. St. Pétersburg., 2: 81—257.
- MOTSCHULSKY V. 1861. Études Entomologiques, Dresden, 10, 24 pp.
- MOTSCHULSKY V. 1866. Catalogue des Insectes reçus du Japon. Bull. Soc. imp. Nat. Mosc., 163—200.
- MBROCZKOWSKI M. 1972. Field investigations in the Democratic People's Republic of Korea by staff members of the Institute of Zoology of the Polish Academy of Sciences. Fragma. faun., 18 (17): 313—343.
- MURAYAMA J. 1934. Jl. Chosen. Nat. Hist. Soc., 19.
- MURAYAMA J. 1937. Jl. Chosen Nat. Hist. Soc., 22.
- MURAYAMA J. 1938. Revision des Sericines (Coléoptères, Scarabéidés) de la Corée. Annot. zool. jap., 17 (1): 7—20.
- MURAYAMA J. 1941. Nouvelles Espèces de Scarabaeides du Manchoukuo et de la Corée. Annot. zool. jap., 20 (1): 36—40.
- MURAYAMA J. 1954. Icones of the Scarabaeid-Beetles from Manchuria and Korea. I, 163 pp.
- NAKANE T. 1951. New or little known *Coleoptera* from Japan and its adjacent Regions. VI. Coprophagous *Lamellicornia*. Ent. Rev. Japan, 5: 69—72.
- NAKANE T. 1956. Neune Aphodiinen aus Japan. Insecta matsum., 20: 119—124.
- NAKANE T. 1960. On the *Scarabaeidae* of Japan (VI). Ent. Rev. Japan, 12 (1): 20—33.
- NAKANE T. 1961. On the *Scarabaeidae* of Japan (VII). Ent. Rev. Japan, 12 (2): 50—63.
- NAKANE T. 1973. Notes on *Serica higonina* LEWIS (*Coleoptera*, *Scarabaeidae*). Ent. Rev. Japan, 25 (1—2): 63—64.
- NAKANE T. 1977. Coloured illustrations of the insects of Japan (*Coleoptera*), I. Osaka, 274 pp.
- NAKANE T., BABA K. 1960. *Scarabaeidae* of Niigata Pref., Japan. In: Insects of Niigata Pref., 4: 1—9.
- NAKANE T., MASUMOTO K. 1967. Studies on the Subgenus *Acrossus* MULSANT (Genus *Aphodius* ILLIGER). Ent. Rev. Japan, 19 (1): 35—39.
- NEWMANN E. 1838. New species of *Popilliae* LEACH. Mag. nat. Hist., 2 (2): 336—338.
- NIIJIMA Y., KINOSHITA E. 1923. Die Untersuchungen über Japanische Melolonthiden, II. Res. Bull. Coll. Agr. Hokkaido imp. Univ., 2 (2): 1—253.
- NISHIO Y. 1970. The Abdominal Structure of 3 Lucanids, *Lucanus maculifemoratus* MOTSCHULSKY, *Protopocoilus inclinatus* MOTSCHULSKY and *Nipponodorcus rubrofemoratus* VOLLENHOVEN. Ent. Rev. Japan, 22 (2): 94—100.
- NOMURA S. 1942. Die Coprophagen Scarabaeiden aus Nord-Korea. Mushi, 14 (2): 116—117.
- NOMURA S. 1965. Some New Forms of the *Scarabaeoidea* from the Loochoo Islands, III. Ent. Rev. Japan, 18 (1): 1—9.
- NOMURA S. 1966. Zoogeographic study of the *Scarabaeoidea* of the Loochoo Archipelago, with a list of the known species and subspecies from there. Tōhō — Gakuhō, 15: 66—105.
- NOMURA S. 1967. Some new and remarkable species of the *Coleoptera* from Japan and its adjacent Regions, II. Ent. Rev. Japan, 19 (2): 52—62.
- NOMURA S. 1969. Zoogeographical study of the *Scarabaeoidea* of the Izu Islands, with descriptions of some new subspecies. Ent. Rev. Japan, 21 (2): 71—94.
- NOMURA S. 1970. Notes on some Scarabaeid-beetles from Loochoo and Formosa. Ent. Rev. Japan, 22 (2): 61—72.
- NOMURA S. 1971. Notes on the Japanese *Scarabaeidae*, II. Ent. Rev. Japan, 23 (2): 64—72.
- NOMURA S. 1972. On the genus *Serica* from Japan. Tōhō-Gakuhō, 22: 109—137.
- NOMURA S. 1973. On the *Sericini* of Japan (*Scarabaeoidea*). Tōhō — Gakuhō, 23: 120—152.
- NOMURA S. 1973. Notes on the Japanese *Scarabaeidae*, III. Ent. Rev. Japan, 25 (1—2): 13—20.
- NOMURA S. 1973. Notes on the Coprophagous *Lamellicornia* from Taiwan. Ent. Rev. Japan, 25 (1—2): 37—52.
- NOMURA S. 1974. On the *Sericini* of Taiwan (*Col.*, *Scarabaeidae*). Tōhō — Gakuhō, 24: 81—115.
- NOMURA S. 1976. On the *Sericini* of Japan (*Scarabaeoidea*), II. Tōhō — Gakuhō, 26: 167—205.

- NOMURA S., KOBAYASHI H. 1976. On the Genus *Mimela* from Japan and Taiwan (*Col., Scarabaeidae*). Ent. Rev. Japan. **29** (1—2): 34—42.
- NOMURA S., NAKANE T. 1951. On some new and remarkable species of *Aphodiinae* from Japan and Formosa. Kontyû, **19**: 35—47.
- OLIVIER A. G. 1789. Entomologie, ou Histoire Naturelle des Insectes, avec leur caractères génériques et spécifiques, leur description, leur synonymie et leur figure enluminee. Col. Bd. 1. Gatt. 1—8., 497 pp.
- PALLAS P. S. 1781. Icones Insectorum praesertim Russiae Sibiriaeque peculiarium, quae collegit et descriptionibus illustravit. Icon. Ins. Ross., I, VI+342 pp.
- PASCOE F. P. 1863. On certain additions to the genus *Dicranocephalus*. J. ent., **2**: 23—26.
- PETROVITZ R. 1961. Neue und verkannte *Aphodiinae* aus allen Erdteilen. Ent. Arb. Mus. Georg FREY, **12**: 99—135.
- PETROVITZ R. 1972. *Colobotheus indagator* MANN. und *apicalis* HAR. (*Aphodiinae, Scarabaeidae*). Koleopt. Rdsch., **50**: 31—33.
- POUILLAUDE I. 1914. Le genre *Dicranocephalus* HOPE. Insecta IV. Paris.
- REITTER E. 1892. Bestimmungs-Tabelle der Lucaniden und coprophagen Lamellicornien des palaearktischen Faunengebietes. Verh. naturf. Ver. Brünn, **30**: 141—262; **31**: 1—109.
- REITTER E. 1902. Bestimmung-Tabelle der *Melolonthidae* aus der europäischen und angrenzenden Länder. III Verh. naturf. Ver. Brünn **50**: 93—303.
- REITTER E. 1903. Bestimmung-Tabelle der *Melolonthidae*. IV Teil: *Rutelini, Hoplini* und *Glaphyrini*. Verh. naturf. Ver. Brünn, **51**: 28—158.
- SAUNDERS W. W. 1854. Characters of undescribed *Lucanidae*, collected in China by R. FORTUNE. Trans. R. ent. Soc. Lond., **3**: 45—55.
- SCHMIDT A. 1922. *Aphodiinae*. Das Tierreich. Berlin—Leipzig, **45**: XXXVI+614 pp.
- SCHÖNHERR C. J. 1817. Synonymia Insectorum. Erster Band. *Eleutherata* oder Käfer. Skaba, XI+506 pp.
- SCOPOLI I. A. 1772. Annus V. Historico-Naturalis. Lipsiae, 128 pp.
- SOLOVEV V. V. 1961. Sledy drevnego oledenija i periglacialnych uslovij v južnom Primore. Trudy Vsesojuzn. naučno-issled. Geol. Inst., **64**: 141—148.
- SOLSKY S. M. 1874. Description d'une espèce nouvelle du genre *Aphodius*. Col. Hefte, **12**: 13—14.
- SØRENSEN T. 1948. A Method of Establishing Groups of Equal Amplitude in Plant Sociology Based on Similarity of Species Content and its Application to Analyses of the Vegetation of Danish Commons. Biol. skr. København, **5**: 1—34.
- STEBNICKA Z. 1973. Neue Scarabaeidenarten (*Coleoptera*) aus der Koreanischen Volksdemokratischen Republik. Bull. Acad. pol. Sci. (Sér. Sci. biol.), **21** (2): 117—121.
- STEBNICKA Z. 1967. Klucze do oznaczania owadów Polski (Keys for the Identification of Polish Insects). Part 19: Chrząszcze — *Coleoptera*, No 28 a: Żukowate — *Scarabaeidae laparosticti*. Warszawa, 139 pp.
- STEBNICKA Z. 1977. A Revision of the World Species of the Tribe *Aegialiini* (*Coleoptera, Scarabaeidae, Aphodiinae*). Acta zool. cracov., **22** (11): 397—506.
- STEBNICKA Z. 1978. Klucze do oznaczania owadów Polski (Keys for the Identification of Polish Insects). Part 19: Chrząszcze — *Coleoptera*, No. 28 b: Żukowate — *Scarabaeidae pleurosticti*. Warszawa, 63 pp.
- STEBNICKA Z. 1978. New *Aphodius*-Species and Description of the Larva *Aphodius apicalis* HAR. (*Coleoptera, Scarabaeidae, Aphodiinae*). Bull. Acad. pol. Sci. (Sér. Sci. biol.), **26** (4): 257—261.
- TESAŘ Z. 1959. Beitrag zur Kenntnis der Gattungen *Poecilophilides* KRAATZ und *Anthrachophora* BURM. (*Col., Cetoninae*). Čas. slezsk. Mus. Opavě, A. **8**: 1—10.
- TESAŘ Z. 1968. Příspěvek k poznání brouků Mongolska a Sibiře. Čas. slezsk. Mus. Opavě, A, **17**: 169—172.
- YAWATA S. 1942. Trans. Kanzai ent Soc., **13**, No. 2.

- VOLLENHOVEN S. C. 1865. Sur quelques Lucanides du Muséum Royal d'Histoire Naturelle a Leide. Tijdschr. Ent., 8: 137—156.
- WALLACE A. R. 1876. The geographical distribution of animals with a study of the relations of living and extinct faunas as elucidating the past changes of the earth's surface. London, 1, XXIV+503 pp; 2, XII+607 pp.
- WATERHOUSE C. O. 1875. On the Lamellicorn *Coleoptera* of Japan. Trans. R. ent. Soc. Lond., 1: 71—116.
- WESTWOOD J. O. 1840. Descriptions and figures of some new Indian Insects. In: Royle's Himalaya ent., 7: XL—XLII.
- WESTWOOD J. O. 1874. Thesaurus Entomologicus Oxoniensis, or Illustrations of new, rare and interesting Insects, for the most part contained in the collections presented to the University of Oxford by the Rev. F. W. Hope. Oxford, XXIV+205 pp.

STRESZCZENIE

Autorka przedstawia rewizję systematyczno-faunistyczną obejmującą *Scarabaeoidea* Koreańskiej Republiki Ludowo-Demokratycznej, z uwzględnieniem danych bibliograficznych dotyczących całego Półwyspu Koreańskiego. Praca została zrealizowana na podstawie materiałów zebranych przez uczestników sześciu ekspedycji zoologicznych Polskiej Akademii Nauk, które w latach 1959—1974 prowadziły badania w Koreańskiej Republice Ludowo-Demokratycznej. Duża część okazów została zebrana osobiście przez autorkę podczas szóstej „krakowskiej” ekspedycji w 1974 roku. Część szczegółowa opracowania zawiera systematyczny przegląd gatunków, wzbogacony rysunkami aparatów kopulacyjnych, oraz redescription gatunków mało znanych i rzadkich. W omówieniu poszczególnych gatunków zamieszczono odnośne pozycje literatury systematycznej, ważniejsze pozycje literatury faunistycznej, dane o bionomii i ogólnym rozsiedleniu oraz własne komentarze autorki i stanowiska zbadanych okazów. Z ogólnej liczby 224 gatunków wykazanych z Półwyspu Koreańskiego, omówiono w niniejszym opracowaniu 138, w tym 29 nowych dla Korei i jeden nowy dla nauki. Wszystkie gatunki z uwzględnieniem ich rozsiedlenia zestawione są w tabeli. Stanowiska z Korei wykazane w tekście (łącznie ze stanowiskami podawanymi w cytowanej literaturze) zostały oznaczone liczbami i naniesione na mapę schematyczną Półwyspu Koreańskiego według załączonego indeksu. Nazwy geograficzne, podawane w literaturze w języku japońskim zostały przez autorkę transkrybowane w alfabecie łacińskim przy użyciu transliteracji międzynarodowej; pisownia większości tych nazw, wymawianych w języku japońskim, została włączona do indeksu.

Całość opracowania uzupełnia szczegółowa synteza zoogeograficzna, dokonana w oparciu o metodę statystyczną, obejmująca *Scarabaeoidea* Dalekiego Wschodu i nawiązująca do historii trzeciorzędu i czwartorzędu tego obszaru.

